

PYRIMIDINONE COMPOUNDS

The present invention relates to certain novel pyrimidinone compounds, processes for their preparation, intermediates useful in their preparation, pharmaceutical compositions containing them and their use in therapy, in particular in the treatment of atherosclerosis.

WO 95/00649 (SmithKline Beecham plc) describe the phospholipase A2 enzyme Lipoprotein Associated Phospholipase A₂ (Lp-PLA₂), the sequence, isolation and purification thereof, isolated nucleic acids encoding the enzyme, and recombinant host cells transformed with DNA encoding the enzyme. Suggested therapeutic uses for inhibitors of the enzyme included atherosclerosis, diabetes, rheumatoid arthritis, stroke, myocardial infarction, reperfusion injury and acute and chronic inflammation. A subsequent publication from the same group further describes this enzyme (Tew D *et al*, *Arterioscler Thromb Vas Biol* 1996;16;591-9) wherein it is referred to as LDL-PLA₂. A later patent application (WO 95/09921, Icos Corporation) and a related publication in *Nature* (Tjoelker *et al*, vol 374, 6 April 1995, 549) describe the enzyme PAF-AH which has essentially the same sequence as Lp-PLA₂ and suggest that it may have potential as a therapeutic protein for regulating pathological inflammatory events.

It has been shown that Lp-PLA₂ is responsible for the conversion of phosphatidylcholine to lysophosphatidylcholine, during the conversion of low density lipoprotein (LDL) to its oxidised form. The enzyme is known to hydrolyse the sn-2 ester of the oxidised phosphatidylcholine to give lysophosphatidylcholine and an oxidatively modified fatty acid. Both products of Lp-PLA₂ action are biologically active with lysophosphatidylcholine, a component of oxidised LDL, known to be a potent chemoattractant for circulating monocytes. As such, lysophosphatidylcholine is thought play a significant role in atherosclerosis by being responsible for the accumulation of cells loaded with cholesterol ester in the arteries. Inhibition of the Lp-PLA₂ enzyme would therefore be expected to stop the build up of these macrophage enriched lesions (by inhibition of the formation of lysophosphatidylcholine and oxidised free fatty acids) and so be useful in the treatment of atherosclerosis.

The increased lysophosphatidylcholine content of oxidatively modified LDL is also thought to be responsible for the endothelial dysfunction observed in patients with atherosclerosis. Inhibitors of Lp-PLA₂ could therefore prove beneficial in the treatment of this phenomenon. An Lp-PLA₂ inhibitor could also find utility in other disease states that exhibit endothelial dysfunction including diabetes, hypertension, angina pectoris and after ischaemia and reperfusion.

In addition, Lp-PLA₂ inhibitors may also have a general application in any disorder that involves activated monocytes, macrophages or lymphocytes, as all of these cell types express Lp-PLA₂. Examples of such disorders include psoriasis.

Furthermore, Lp-PLA₂ inhibitors may also have a general application in any disorder that involves lipid oxidation in conjunction with Lp-PLA₂ activity to produce the two injurious products, lysophosphatidylcholine and oxidatively modified fatty acids. Such conditions include the


aforementioned conditions atherosclerosis, diabetes, rheumatoid arthritis, stroke, myocardial infarction, reperfusion injury and acute and chronic inflammation. Further such conditions include various neuropsychiatric disorders such as schizophrenia (see Psychopharmacology Bulletin, 31, 159-165, 1995).

Patent applications WO 96/12963, WO 96/13484, WO96/19451, WO 97/02242, WO97/217675, WO97/217676, WO 96/41098, and WO97/41099 (SmithKline Beecham plc) disclose *inter alia* various series of 4-thionyl/sulfinyl/sulfonyl azetidinone compounds which are inhibitors of the enzyme Lp-PLA₂. These are irreversible, acylating inhibitors (Tew *et al*, Biochemistry, 37, 10087, 1998).

Patent applications WO 99/24420 and WO 00/10980 (SmithKline Beecham plc, published after the priority date of the present application) describe a new class of reversible, non-acylating inhibitors of the enzyme Lp-PLA₂, in particular a class of pyrimidone compounds. The early 2-(alkylthio)pyrimidin-4-one chemical lead is described in Bioorganic and Medicinal Chemistry Letters, 2000, 10, 395-8.

A further class of pyrimidone compounds has now been identified which are inhibitors of the enzyme Lp-PLA₂.

Accordingly, the present invention provides a compound of formula (I):



in which:

R¹ is an aryl or heteroaryl group, optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from C₍₁₋₁₈₎alkyl, C₍₁₋₁₈₎alkoxy, C₍₁₋₁₈₎alkylthio, arylC₍₁₋₁₈₎alkoxy, hydroxy, halogen, CN, COR⁶, carboxy, COOR⁶, CONR⁹R¹⁰, NR⁶COR⁷, SO₂NR⁹R¹⁰, NR⁶SO₂R⁷, NR⁹R¹⁰, mono to perfluoro-C₍₁₋₄₎alkyl and mono to perfluoro-C₍₁₋₄₎alkoxy, or, as a single substituent, optionally in combination with a further substituent as hereinbefore defined, CH₂COOH or a salt thereof, CH₂COOR⁸, CH₂CONR⁹R¹⁰, CH₂CN, (CH₂)_mNR⁹R¹⁰, (CH₂)_mOH or (CH₂)_mOR⁶ where m is an integer from 1 to 3;

R² is an aryl or heteroaryl group, optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from C₍₁₋₁₈₎alkyl, C₍₁₋₁₈₎alkoxy, C₍₁₋₁₈₎alkylthio, arylC₍₁₋₁₈₎alkoxy, hydroxy, halogen, CN, COR⁶, carboxy, COOR⁶, CONR⁹R¹⁰, NR⁶COR⁷, SO₂NR⁹R¹⁰, NR⁶SO₂R⁷, NR⁹R¹⁰, mono to perfluoro-C₍₁₋₄₎alkyl, mono to perfluoro-C₍₁₋₄₎alkoxy, and arylC₍₁₋₄₎alkyl;

R^3 is hydrogen or $C_{(1-4)}$ alkyl which may be unsubstituted or substituted by hydroxy, OR^6 , COR^6 , carboxy, $COOR^6$, $CONR^9R^{10}$, NR^9R^{10} , mono- or di-(hydroxy $C_{(1-6)}$ alkyl)amino or N-hydroxy $C_{(1-6)}$ alkyl-N- $C_{(1-6)}$ alkyl amino;

5 R^4 is an aryl or a heteroaryl ring optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from $C_{(1-18)}$ alkyl, $C_{(1-18)}$ alkoxy, $C_{(1-18)}$ alkylthio, aryl $C_{(1-18)}$ alkoxy, hydroxy, halogen, CN, COR^6 , carboxy, $COOR^6$, $CONR^9R^{10}$, NR^6COR^7 , $SO_2NR^9R^{10}$, $NR^6SO_2R^7$, NR^9R^{10} , mono to perfluoro- $C_{(1-4)}$ alkyl and mono to perfluoro- $C_{(1-4)}$ alkoxy;

10 R^5 is an aryl ring which is further optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from $C_{(1-18)}$ alkyl, $C_{(1-18)}$ alkoxy, $C_{(1-18)}$ alkylthio, aryl $C_{(1-18)}$ alkoxy, hydroxy, halogen, CN, COR^6 , carboxy, $COOR^6$, $CONR^9R^{10}$, NR^6COR^7 , $SO_2NR^9R^{10}$, $NR^6SO_2R^7$, NR^9R^{10} , mono to perfluoro- $C_{(1-4)}$ alkyl and mono to perfluoro- $C_{(1-4)}$ alkoxy;

R^6 and R^7 are independently hydrogen or $C_{(1-20)}$ alkyl, for instance $C_{(1-4)}$ alkyl (e.g. methyl or ethyl);

R^8 is $C_{(1-4)}$ alkyl or a pharmaceutically acceptable *in vivo* hydrolysable ester group;

15 R^9 and R^{10} which may be the same or different is each selected from hydrogen, $C_{(1-12)}$ alkyl, CH_2R^{11} , $CHR^{12}CO_2H$ or a salt thereof, or R^9 and R^{10} together with the nitrogen to which they are attached form a 4- to 7-, preferably 5- to 7-, membered ring optionally containing one or more further heteroatoms selected from oxygen, nitrogen and sulphur, and optionally substituted by one or two substituents selected from hydroxy, oxo, $C_{(1-4)}$ alkyl, $C_{(1-4)}$ alkylCO, aryl, e.g. phenyl, or aralkyl, e.g. benzyl, for instance morpholine or piperazine;

R^{11} is $COOH$ or a salt thereof, $COOR^8$, $CONR^6R^7$, CN, CH_2OH or CH_2OR^6 ;

R^{12} is an amino acid side chain such as CH_2OH from serine;

n is an integer from 1 to 4, preferably 1 or 3;

X is O or S; and

25 Z is $CR^{13}R^{14}$ where R^{13} and R^{14} are each hydrogen or $C_{(1-4)}$ alkyl, or R^{13} and R^{14} together with the intervening carbon atom form a $C_{(3-6)}$ cycloalkyl ring.

Preferably, Z is CH_2 .

30 Representative examples of R^1 when an aryl group include phenyl and naphthyl.

Representative examples of R^1 when a heteroaryl group include a 5- or 6- membered, monocyclic heteroaryl group comprising 1 or 2 nitrogen heteroatoms.

35 Preferably, R^1 is pyrimidyl optionally substituted by 1 or 2 substituents preferably selected from oxo, aryl $C_{(1-4)}$ alkyl (e.g. benzyl), $C_{(1-6)}$ alkyl (e.g. methyl or ethyl), $C_{(3-6)}$ cycloalkyl, hydroxy, $C_{(1-4)}$ alkoxy (e.g. methoxy), carboxy $C_{(1-6)}$ alkyl, $C_{(1-6)}$ alkylcarboxy $C_{(1-6)}$ alkyl, di- $C_{(1-6)}$ alkylamino, and morpholino; or pyrazolyl optionally substituted by $C_{(1-6)}$ alkyl (e.g. methyl or ethyl).

40 Preferably, ZR^1 is pyrimid-5-ylmethyl optionally substituted by 2-methoxy, 2-trifluoromethyl, 2-(4-morpholino) or 2-dimethylamino; 2-oxo-pyrimid-5-ylmethyl or 1-methyl-4-pyrazolylmethyl.

Preferably X is S.

Representative examples of R^2 when an aryl group include phenyl and naphthyl. Representative examples of R^2 when a heteroaryl group include pyridyl, pyrimidinyl, pyrazolyl, furanyl, thienyl, thiazolyl, quinolyl, benzothiazolyl, pyridazolyl and pyrazinyl. Preferably, R^2 is phenyl optionally substituted by halogen.

Representative examples of R^2CH_2X include 4-fluorobenzylthio.

Representative examples of R^3 include hydrogen; and methyl, ethyl and propyl, optionally substituted by amino, $C_{(1-3)}$ alkylamino, di $C_{(1-3)}$ alkyl amino, hydroxy $C_{(1-3)}$ alkylamino, hydroxy, $C_{(1-3)}$ alkoxy, carboxy, $C_{(1-3)}$ alkylcarboxy, and heterocycyl such as piperidino, piperazino, pyrrolidono and morpholino, wherein the alkyl moiety, if present, is preferably methyl, or ethyl.

Representative examples of R^4 include phenyl optionally substituted by halogen; thiophene; pyridine; and pyrimidine.

Representative examples of R^5 include phenyl optionally substituted by halogen, trifluoromethyl, or trifluoromethoxy, preferably at the 4-position.

Preferably, R^4 and R^5 together form a 4-(phenyl)phenyl substituent in which the remote phenyl ring may be optionally substituted by halogen or trifluoromethyl, preferably at the 4-position.

Pharmaceutically acceptable *in vivo* hydrolysable ester groups for R^8 include those which break down readily in the human body to leave the parent acid or its salt.

Representative examples of values of pharmaceutically acceptable *in vivo* hydrolysable ester groups for R^8 include:


$-CH(R^a)O.CO.R^b$;

$-CH(R^a)O.CO.OR^c$;


$-CH(R^a)CO.NR^eR^f$

$-R^dNR^eR^f$;

$-CH_2OR^g$;



$CH(R^a)O.CO.C_6H_4Y^1COCH(R^i)NH_2$; and



in which:

R^a is hydrogen, (C₁₋₆)alkyl, in particular methyl, (C₃₋₇)cycloalkyl, or phenyl, each of which may be optionally substituted;

R^b is (C₁₋₆)alkyl, (C₁₋₆)alkoxy(C₁₋₆)alkyl, phenyl, benzyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl(C₃₋₇)cycloalkyl, 1-amino(C₁₋₆)alkyl, or 1-(C₁₋₆alkyl)amino(C₁₋₆)alkyl, each of which may be optionally substituted; or

R^a and R^b together form a 1,2-phenylene group optionally substituted by one or two methoxy groups;

R^c is (C₁₋₆)alkyl, (C₃₋₇)cycloalkyl, (C₁₋₆)alkyl(C₃₋₇)cycloalkyl;

R^d is (C₁₋₆)alkylene optionally substituted with a methyl or ethyl group;

R^e and R^f which may be the same or different is each (C₁₋₆)alkyl; or aryl(C₁₋₄) alkyl, optionally substituted with e.g. hydroxy;

R^g is (C₁₋₆)alkyl;

R^h is hydrogen, (C₁₋₆)alkyl or phenyl;

Rⁱ is hydrogen or phenyl optionally substituted by up to three groups selected from halogen, (C₁₋₆)-alkyl, or (C₁₋₆)alkoxy;

and

Y¹ is oxygen or NH;

for instance:

(a) acyloxyalkyl groups such as acetoxymethyl, isobutyryloxymethyl, pivaloyloxymethyl, benzoyloxymethyl, α -acetoxylethyl, α -pivaloyloxyethyl, 1-(cyclohexylcarbonyloxy)ethyl, (1-aminoethyl)carbonyloxymethyl, 2-methoxyprop-2-ylcarbonyloxymethyl, phenylcarbonyloxymethyl and 4-methoxyphenyl-carbonyloxymethyl;

(b) alkoxy/cycloalkoxycarbonyloxyalkyl groups, such as ethoxycarbonyloxymethyl, t-butyloxycarbonyloxymethyl, cyclohexyloxycarbonyloxymethyl, 1-methylcyclohexyloxycarbonyloxymethyl and α -ethoxycarbonyloxyethyl;

(c) dialkylaminoalkyl, especially di-loweralkylamino alkyl groups such as dimethylaminomethyl, dimethylaminoethyl, diethylaminomethyl or diethylaminoethyl;

(d) acetamido groups such as N,N-dimethylaminocarbonylmethyl, N,N-(2-hydroxyethyl)aminocarbonylmethyl;

(e) lactone groups such as phthalidyl and dimethoxyphthalidyl;

(f) (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl; and

(g) (2-methoxycarbonyl-E-but-2-en-yl)methyl.

Representative examples of pharmaceutically acceptable *in vivo* hydrolysable ester groups for R⁸ include:

(2-methoxycarbonyl-E-but-2-en-yl)methyl, isobutyryloxymethyl, 2-methoxyprop-2-ylcarbonyloxymethyl, phenylcarbonyloxymethyl, 4-methoxyphenyl-carbonyloxymethyl, t-butyloxycarbonyloxymethyl, cyclohexyloxy-carbonyloxymethyl, 1-methylcyclohexyloxycarbonyloxymethyl, N,N-dimethylaminocarbonylmethyl, and (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl.

It will be appreciated that in some instances, compounds of the present invention may include a basic function such as an amino group as a substituent. Such basic functions may be used to form acid addition salts, in particular pharmaceutically acceptable salts. Pharmaceutically acceptable salts

include those described by Berge, Bighley, and Monkhouse, *J. Pharm. Sci.*, 1977, 66, 1-19. Such salts may be formed from inorganic and organic acids. Representative examples thereof include maleic, fumaric, benzoic, ascorbic, pamoic, succinic, bismethylenesalicylic, methanesulfonic, p-toluenesulfonic, ethanedisulfonic, acetic, propionic, tartaric, salicylic, citric, gluconic, aspartic, stearic, 5 palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, taurocholic acid, benzenesulfonic, hydrochloric, hydrobromic, sulfuric, cyclohexylsulfamic, phosphoric and nitric acids.

It will be appreciated that in some instances, compounds of the present invention may include a carboxy group as a substituent. Such carboxy groups may be used to form salts, in particular pharmaceutically acceptable salts. Pharmaceutically acceptable salts include those described by Berge, Bighley, and 10 Monkhouse, *J. Pharm. Sci.*, 1977, 66, 1-19. Preferred salts include alkali metal salts such as the sodium and potassium salts.

When used herein, the term "alkyl" and similar terms such as "alkoxy" includes all straight chain and 15 branched isomers. Representative examples thereof include methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *t*-butyl, *n*-pentyl and *n*-hexyl.

When used herein, the term "aryl" refers to, unless otherwise defined, a mono- or bicyclic aromatic ring system containing up to 10 carbon atoms in the ring system, for instance phenyl or naphthyl. 20

When used herein, the term "heteroaryl" refers to a mono- or bicyclic heteroaromatic ring system comprising up to four, preferably 1 or 2, heteroatoms each selected from oxygen, nitrogen and sulphur. Each ring may have from 4 to 7, preferably 5 or 6, ring atoms. A bicyclic heteroaromatic ring system may include a carbocyclic ring. 25

When used herein, the terms "halogen" and "halo" include fluorine, chlorine, bromine and iodine and fluoro, chloro, bromo and iodo, respectively.

Particularly preferred compounds of formula (I) are:

30 1-(N-Methyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methylpyrazol-4-ylmethyl)pyrimidin-4-one;
1-(N-Methyl-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methylpyrazol-4-ylmethyl)pyrimidin-4-one;
1-(N-(2-Dimethylaminoethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluoro- 35 benzyl)thio-5-(1-methylpyrazol-4-ylmethyl)pyrimidin-4-one;
1-(N-Methyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-(4-morpholino)pyrimidin-5-ylmethyl)pyrimidin-4-one;
1-(N-(2-(dimethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one;
40 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one;
1-(N-(2-(diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyridin-5-ylmethyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one;

1-(N-(2-(1-Piperidino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one bitartrate;
1-(N-(Carboxymethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one sodium salt; and
5 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one or a pharmaceutically acceptable salt thereof, including the hydrochloride, bitartrate, citrate and tosylate salts.

10 Since the compounds of the present invention, in particular compounds of formula (I), are intended for use in pharmaceutical compositions, it will be understood that they are each provided in substantially pure form, for example at least 50% pure, more suitably at least 75% pure and preferably at least 95% pure (% are on a wt/wt basis). Impure preparations of the compounds of formula (I) may be used for preparing the more pure forms used in the pharmaceutical compositions. Although the purity of intermediate compounds of the present invention is less critical, it will be readily understood that the
15 substantially pure form is preferred as for the compounds of formula (I). Preferably, whenever possible, the compounds of the present invention are obtained in crystalline form.

20 When some of the compounds of this invention are allowed to crystallise or are recrystallised from organic solvents, solvent of crystallisation may be present in the crystalline product. This invention includes within its scope such solvates. Similarly, some of the compounds of this invention may be crystallised or recrystallised from solvents containing water. In such cases water of hydration may be formed. This invention includes within its scope stoichiometric hydrates as well as compounds containing variable amounts of water that may be produced by processes such as lyophilisation. In addition, different crystallisation conditions may lead to the formation of different polymorphic forms
25 of crystalline products. This invention includes within its scope all polymorphic forms of the compounds of formula (I).

30 Compounds of the present invention are inhibitors of the enzyme lipoprotein associated phospholipase A₂ (Lp-PLA₂) and as such are expected to be of use in therapy, in particular in the treatment of atherosclerosis. In a further aspect therefore the present invention provides a compound of formula (I) for use in therapy.

35 The compounds of formula (I) are inhibitors of lysophosphatidylcholine production by Lp-PLA₂ and may therefore also have a general application in any disorder that involves endothelial dysfunction, for example atherosclerosis, diabetes, hypertension, angina pectoris and after ischaemia and reperfusion. In addition, compounds of formula (I) may have a general application in any disorder that involves lipid oxidation in conjunction with enzyme activity, for example in addition to conditions such as atherosclerosis and diabetes, other conditions such as rheumatoid arthritis, stroke, inflammatory conditions of the brain such as Alzheimer's Disease, myocardial infarction, reperfusion injury, sepsis,
40 and acute and chronic inflammation. Further such conditions include various neuropsychiatric disorders such as schizophrenia (see Psychopharmacology Bulletin, 31, 159-165, 1995).

Further applications include any disorder that involves activated monocytes, macrophages or lymphocytes, as all of these cell types express Lp-PLA₂. Examples of such disorders include psoriasis.

Accordingly, in a further aspect, the present invention provides for a method of treating a disease state associated with activity of the enzyme Lp-PLA₂ which method involves treating a patient in need thereof with a therapeutically effective amount of an inhibitor of the enzyme. The disease state may be associated with the increased involvement of monocytes, macrophages or lymphocytes; with the formation of lysophosphatidylcholine and oxidised free fatty acids; with lipid oxidation in conjunction with Lp PLA₂ activity; or with endothelial dysfunction.

Compounds of the present invention may also be of use in treating the above mentioned disease states in combination with an anti-hyperlipidaemic, anti-atherosclerotic, anti-diabetic, anti-anginal, anti-inflammatory, or anti-hypertension agent or an agent for lowering Lp(a). Examples of the above include cholesterol synthesis inhibitors such as statins, anti-oxidants such as probucol, insulin sensitisers, calcium channel antagonists, and anti-inflammatory drugs such as NSAIDs. Examples of agents for lowering Lp(a) include the aminophosphonates described in WO 97/02037, WO 98/28310, WO 98/28311 and WO 98/28312 (Symphar SA and SmithKline Beecham).

A preferred combination therapy will be the use of a compound of the present invention and a statin. The statins are a well known class of cholesterol lowering agents and include atorvastatin, simvastatin, pravastatin, cerivastatin, fluvastatin, lovastatin and ZD 4522 (also referred to as S-4522, Astra Zeneca). The two agents may be administered at substantially the same time or at different times, according to the discretion of the physician.

A further preferred combination therapy will be the use of a compound of the present invention and an anti-diabetic agent or an insulin sensitiser, as coronary heart disease is a major cause of death for diabetics. Within this class, preferred compounds for use with a compound of the present invention include the PPAR γ activators, for instance G1262570 (Glaxo Wellcome) and also the glitazone class of compounds such as rosiglitazone (Avandia, SmithKline Beecham), troglitazone and pioglitazone.

In therapeutic use, the compounds of the present invention are usually administered in a standard pharmaceutical composition. The present invention therefore provides, in a further aspect, a pharmaceutical composition comprising a compound of formula (I) and a pharmaceutically acceptable carrier.

Suitable pharmaceutical compositions include those which are adapted for oral or parenteral administration or as a suppository.


Suitable pharmaceutical compositions include those which are adapted for oral or parenteral administration or as a suppository. Compounds of formula (I) which are active when given orally can be formulated as liquids, for example syrups, suspensions or emulsions, tablets, capsules and lozenges. A liquid formulation will generally consist of a suspension or solution of the compound or

pharmaceutically acceptable salt in a suitable liquid carrier(s) for example, ethanol, glycerine, non-aqueous solvent, for example polyethylene glycol, oils, or water with a suspending agent, preservative, flavouring or colouring agent. A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose. A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule. Typical parenteral compositions consist of a solution or suspension of the compound of formula (I) in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration. A typical suppository formulation comprises a compound of formula (I) which is active when administered in this way, with a binding and/or lubricating agent such as polymeric glycols, gelatins or cocoa butter or other low melting vegetable or synthetic waxes or fats.

Preferably the composition is in unit dose form such as a tablet or capsule. Each dosage unit for oral administration contains preferably from 1 to 500 mg (and for parenteral administration contains preferably from 0.1 to 25 mg) of a compound of the formula (I). The daily dosage regimen for an adult patient may be, for example, an oral dose of between 1 mg and 1000 mg, preferably between 1 mg and 500 mg, or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 25 mg, of the compound of the formula (I), the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

A compound of formula (I) may be prepared by a number of processes which include:

(a) reacting a compound of formula (II):



(II)


in which X, Y, Z, R¹ and R² are as hereinbefore defined, with a compound of formula (III):



(III)


in which R³, R⁴ and R⁵ are as hereinbefore defined; under amide forming conditions;

(b) reacting a compound of formula (IV):



(IV)


- 5 in which X, Z, R¹ and R² are as hereinbefore defined,
with a compound of formula (V):



(V)

- 10 in which n, R³, R⁴ and R⁵ are as hereinbefore defined, and L¹ is a leaving group such as halogen, for instance bromo or iodo,
in the presence of a base such as a secondary or tertiary amine, for instance di-iso-propylethylamine, in an inert solvent such as dichloromethane;

- 15 (c) when X is S, reacting a compound of formula (VI):



(VI)


- 20 in which n, Z, R¹, R³, R⁴ and R⁵ are as hereinbefore defined,
with a compound of formula (VII):



(VII)

- 25 in which R² and L¹ are as hereinbefore defined,
in the presence of a base such as a secondary or tertiary amine, for instance di-isopropylethylamine, in an inert solvent such as dichloromethane; or

(d) when X is O, reacting a compound of formula (VIII):



(VIII)

in which n , Z , R^1 , R^3 , R^4 and R^5 are as hereinbefore defined, and L^2 is a leaving group such as halogen or alkylthio, for instance methylthio,

5 with a compound of formula (IX):




(IX)

in which R^2 is as hereinbefore defined,

10 in the presence of a base such as 4-dimethylaminopyridine, in an inert solvent such as pyridine.

Compounds of formulae (II), (IV), (VI) and (VIII) for use in the above processes may be prepared by processes illustrated in the following scheme I:



in which:

L^3 is a C(1-6)alkyl group, for instance methyl;

- 5 R^{15} is a C(1-6)alkyl group, for instance methyl, ethyl, or t-butyl, and $L^1, L^2, R^1, R^2, R^3, R^4, R^5, n, X$ and Z are as hereinbefore defined.

With reference to Scheme I:

- 10 Amide forming conditions for step (a) are well known in the art. Preferably, the acid of formula (II) is reacted with the amine of formula (III) in an inert solvent, such as dichloromethane, at ambient temperature and in the presence of an activating agent such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide plus hydroxybenzotriazole.

Alkylation conditions for step (b) include reaction in the presence of a base such as a secondary or tertiary amine, for instance di-*iso*-propylethylamine, in an inert solvent such as dichloromethane, forming an intermediate ester which is converted to the acid of formula (II) by hydrolysis, for instance using aqueous sodium hydroxide in a solvent such as dioxan or by alternative deprotection, for instance
5 using trifluoroacetic acid in a solvent such as dichloromethane.

Conditions for step (c) include under thioether forming conditions. Advantageously, the reaction is carried out in the presence of a base such as sodium ethoxide or potassium carbonate, preferably in a solvent such as ethanol or dimethyl formamide, or a secondary or tertiary amine base such as di-
10 isopropylethyl amine, in solvent such as dichloromethane.

In step (d), a compound of formula (XVII) is reacted with thiourea, in the presence of sodium ethoxide (preferably generated *in situ* from sodium and ethanol).

15 In step (e), a compound of formula (XVIII) is reacted with ethyl formate in the presence of a base such as sodium hydride or potassium isopropoxide.

In step (f), a compound of formula (IV) is reacted with a compound of formula (V) in the presence of a base such as a secondary or tertiary amine, for instance di-isopropylethylamine, in an inert solvent such
20 as dichloromethane

In step (g), a compound of formula (XIII) is reacted with a compound of formula (XIV) in a solvent such as dimethylformamide to form an intermediate thiourea, which is then treated with a base such as sodium methoxide.
25

In step (h), a compound of formula (XVI) is reacted with a metal thiocyanate, for example potassium thiocyanate, in a solvent such as acetonitrile.

30 In step (i), a compound of formula (XVII) is reacted with a methylating agent such as dimethyl sulphate in the presence of a base such as potassium carbonate, followed by hydrolysis of the intermediate ester in conventional manner e.g. by basic hydrolysis using sodium hydroxide to give the corresponding carboxylic acid which may then be converted into the acyl chloride, for instance by treatment with oxalyl chloride.

35 In step (j), a catalyst such as 4-dimethylaminopyridine, and in a solvent such as pyridine are used.


In step (k), a compound of formula (XIII) is reacted with a compound of formula (XV) in a solvent such as dimethylformamide to form an intermediate thiourea, which is then treated with a base such as sodium methoxide.
40

The present invention will now be illustrated by the following examples.

Examples

The structure and purity of the intermediates and examples was confirmed by ¹H-NMR and (in nearly all cases) mass spectroscopy, even where not explicitly indicated below.


5 Intermediate A1 — 4-(4-Chlorophenyl)benzaldehyde



(a) A mixture of 4-formylbenzeneboronic acid (2.50g, 2 equiv), 4-chloriodobenzene (1.98g, 1 equiv), tetrakis(triphenylphosphine)palladium(0) (0.50g, 0.05 equiv), aqueous sodium carbonate (18ml, 2M solution, 2 equiv) and dimethoxyethane (50ml) was stirred at reflux under argon overnight, then cooled and diluted with ethyl acetate. The mixture was filtered as necessary to remove inorganic residues, then the organic layer was washed successively with aqueous citric acid and brine, dried and evaporated. The crude product was purified by chromatography (silica, 5% ethyl acetate in hexane); product fractions were evaporated to a white solid (1.32g, 72%).


(b) A mixture of 4-chlorobenzeneboronic acid (19.4g, 1 equiv), 4-bromobenzaldehyde (22.9g, 1 equiv), palladium(II) acetate (1.4g, 0.05 equiv) aqueous sodium carbonate (30.3 g in 144ml solution, 2 equiv) and dimethoxyethane (500ml) was stirred at reflux under argon for 2.5h, then evaporated to low volume and diluted with dichloromethane. Workup continued as in (a) above to give identical material (25.2g, 94%). ¹H-NMR (CDCl₃) δ 10.05 (1H, s), 7.96 (2H, d), 7.73 (2H, d), 7.57 (2H, d), 7.46 (2H, d); MS (AP⁺) found (M+1) = 217, C₁₃H₉³⁵ClO requires 216.

20 Intermediate A2 — N-Methyl-4-(4-chlorophenyl)benzylamine



A mixture of Intermediate A1 (3.5g, 1 equiv), methylamine (32.3ml of a 2M solution in THF, 4 equiv) and anhydrous magnesium sulphate (4.47g, 2 equiv) was stirred at room temperature for 16h, then filtered, the solid washed thoroughly with ethyl acetate, and the combined filtrates evaporated to a white solid (3.7g). This imine intermediate was suspended in ethanol (100ml), cooled in ice and sodium borohydride (0.61g, 1 equiv) added portionwise. The ice bath was removed, and the mixture stirred for 45min at room temperature then at 50°C for 1h. The solvent was removed in vacuo, water was added to the residue, and the product extracted into dichloromethane. Drying and evaporation of the solvent gave a white solid (3.56g). ¹H-NMR (CDCl₃) δ 7.51 (4H, d), 7.40 (4H, d), 3.79 (2H, s), 2.48 (3H, s); MS (APCI⁺) found (M+1) = 232, C₁₄H₁₄³⁵ClN requires 231.


Intermediate A3 — N-(2-Diethylaminoethyl)-4-(4-chlorophenyl)benzylamine



A mixture of Intermediate A1 (55.0g), N,N-diethylethylenediamine (35.6ml), 4A molecular sieve (37g), and dichloromethane (1100ml) was reacted at room temperature under argon for 16h, with occasional agitation. The solid was filtered off and washed with dichloromethane, and the combined filtrates evaporated to a yellow foam (72.4g). This intermediate imine was reduced with sodium borohydride


(8.7g) in ethanol (850ml) as described for Intermediate A2, yielding the title compound as a yellow oil (72.7g). $^1\text{H-NMR}$ (CDCl_3) δ 1.70 (2H, t), 2.22 (6H, s), 2.33 (2H, t), 2.69 (2H, br. m), 3.83 (2H, s), 7.37-7.43 (4H, m), 7.52-7.56 (4H, m).

Intermediate A4 — 4-(4-Chlorophenyl)benzyl alcohol




A mixture of Intermediate A1 (1.73g, 1 equiv), sodium borohydride (0.3g, 1 equiv) and ethanol (20ml) was stirred until a clear solution was obtained; a gentle exotherm was observed. The temperature was raised to 50°C and stirring continued for 2h, then the solvent was removed in vacuo, water added to the residue, and the product extracted into dichloromethane. Drying and evaporation of the solvent gave a white solid (1.67g). $^1\text{H-NMR}$ (CDCl_3) δ 1.73 (1H, t), 4.74 (2H, d), 7.2-7.6 (8H, m).

Intermediate A5 — N-(4-(4-Chlorophenyl)benzyl)phthalimide




Intermediate A4 (1.62g, 1 equiv), phthalimide (1.42g, 1.3 equiv) and triphenylphosphine (2.53g, 1.3 equiv) were dissolved in dry THF (50ml), the solution was cooled in ice, and diethyl azodicarboxylate (1.52ml, 1.3 equiv) was added slowly with stirring. The yellow solution was stirred for 16h at room temperature, then the solvent was evaporated. Chromatography (silica, 1:1 dichloromethane/pet. ether) and trituration with ether gave the desired product (2.03g, 79%). $^1\text{H-NMR}$ (CDCl_3) δ 4.89 (2H, s), 7.36-7.50 (8H, m), 7.69-7.74 (2H, m), 7.82-7.89 (2H, m).

Intermediate A6 — 4-(4-Chlorophenyl)benzylamine



A mixture of Intermediate A5 (1.98g, 1 equiv) and hydrazine hydrate (0.55ml, 2 equiv) in ethanol (40ml) was heated at reflux for 2h, then the solvent was evaporated. The residue was shaken with dichloromethane and 0.5M aqueous sodium hydroxide, the aqueous layer re-extracted with dichloromethane, and the combined organic extracts purified by chromatography (silica, 1-3% methanolic ammonia in dichloromethane). Product fractions were evaporated to a white solid (0.53g, 43%). $^1\text{H-NMR}$ (CDCl_3) δ 3.92 (2H, s), 7.28-7.41 (4H, m), 7.49-7.54 (4H, m).


Intermediate A7 — N-Methyl-N-(4-(4-chlorophenyl)benzyl)bromoacetamide



Bromoacetyl bromide (0.123ml, 1.1 equiv) was added to a well-stirred mixture of Intermediate A2 (0.3g, 1 equiv), sodium hydroxide (0.06g, 1.2 equiv), water and dichloromethane. Stirring was continued for 3h at room temperature, then the phases were separated and the organic layer dried and evaporated. Chromatography (silica, 3:1 hexane/ethyl acetate) gave the desired product as an oil


(0.37g). $^1\text{H-NMR}$ (CDCl_3) δ 7.38-7.68 (8H, m), 4.77 (2H, s), 4.07 (2H, s), 3.24 (3H, s); MS (APCI+) found $(\text{M}+1) = 352$, $\text{C}_{16}\text{H}_{15}^{79}\text{Br}^{35}\text{ClNO}$ requires 351.

Intermediate A8 — 5-Hydroxymethyl-2-(4-fluorophenyl)pyridine




- 5 A solution of Intermediate A23 (4.63g) in dry dichloromethane (100ml) was cooled to -78°C under argon, then DIBAL-H (26.7ml, 1.5M solution in toluene) was added dropwise over 20min. Stirring was continued for 40min at -78°C , then 2M hydrochloric acid (52ml) was added dropwise over 15min. The solution was allowed to warm slowly to room temperature, then the organic layer was separated, washed with water, dried and evaporated. Chromatography (silica, 1:1 ethyl acetate/hexane) gave the product as a white solid (3.03g, 75%). $^1\text{H-NMR}$ (CDCl_3) δ (1H, d), 7.98 (2H, m), 7.77 (1H, m), 7.68 (1H, d), 7.15 (2H, t), 4.77 (2H, d), 1.802 (1H, t); MS(APCI+) found $(\text{M}+1)=204$, $\text{C}_{12}\text{H}_{10}\text{FNO}$ requires 203.

Intermediate A9 — 5-Formyl-2-(4-fluorophenyl)pyridine




- 15 Activated manganese dioxide (3.19g) was added to a solution of Intermediate A8 (0.75g) in dichloromethane (50ml) and stirred at room temperature for 16h. The solids were filtered off and the filtrate evaporated to a pale yellow solid (0.57g). $^1\text{H-NMR}$ (CDCl_3) δ 10.15 (1H, s), 9.11 (1H, s), 8.22 (1H, dd), 8.10 (2H, m), 7.87 (1H, d), 7.20 (2H, t); MS(APCI+) found $(\text{M}+1)=202$, $\text{C}_{12}\text{H}_8\text{FNO}$ requires 201.

Intermediate A10 — N-Methyl-4-(4-chlorophenyl)-3-fluorobenzylamine



- Borane (1M solution in THF, 2 equiv) was stirred under argon with ice cooling, and a suspension of Intermediate A39 (1 equiv) in dry THF was added gradually over a few minutes. After 5 mins the ice bath was removed and the mixture heated to reflux for 1h, then cooled to room temperature. 5M Hydrochloric acid was added dropwise, and the THF was removed from the resulting suspension by distillation. The residue was diluted with water and made strongly basic with sodium hydroxide, then the product was extracted into ether. Drying and evaporation of the organic solution, followed by chromatography gave the title compound as an oil (88%). $^1\text{H-NMR}$ (CDCl_3) δ 2.48 (3H, s), 3.79(2H, s), 7.12-7.18 (2H, m), 7.33-7.50 (5H, m); MS (APCI+) found $(\text{M}+1) = 250$; $\text{C}_{14}\text{H}_{13}\text{ClFN}$ requires 249.


Intermediate A11 — N-(Ethoxycarbonylmethyl)-4-(4-chlorophenyl)benzylamine



A mixture of Intermediate A1 (0.5 g, 1 equiv), glycine ethyl ester hydrochloride (0.32 g, 1 equiv), diisopropylethylamine (0.4 ml, 1 equiv) and 1,2-dichloroethane (10 ml) was stirred at room temperature, sodium triacetoxyborohydride (0.73 g, 1.5 equiv) was added, and stirring continued

overnight. After diluting with dichloromethane, the solution was washed with water and dried over potassium carbonate. Evaporation of the solvent yielded the title compound (0.57 g) as a white waxy solid. $^1\text{H-NMR}$ (CDCl_3) δ 1.22-1.37 (3H, t), 3.44 (2H, s), 3.85 (2H, s), 4.15-4.32 (2H, q), 7.35-7.62 (8H, m); MS (APCI+) found ($M+1$) = 304; $\text{C}_{17}\text{H}_{18}\text{ClNO}_2$ requires 303.


5 **Intermediate A12 — 2-Hydroxymethyl-5-(4-chlorophenyl)pyridine**



m-Chloroperbenzoic acid (0.93 g) was added portionwise over 30 min to a solution of Intermediate A41 (0.77 g) in dichloromethane (10 ml), then the mixture was stirred at room temperature for a further hour, diluted with dichloromethane, washed with aq. sodium bicarbonate, dried and evaporated.


- 10 Chromatography (silica, 5% methanol in dichloromethane) gave 2-methyl-5-(4-chlorophenyl)pyridine-N-oxide (0.79 g) as a white solid. This material was dissolved in dichloromethane (15 ml), trifluoroacetic anhydride (1.9 ml) added, and the mixture stirred for 1 hour at room temperature followed by 1 hour at reflux. Volatile components were removed in vacuo, then the residue was redissolved in dichloromethane (5 ml), 2M aq. sodium carbonate (14 ml) was added, and stirred
- 15 vigorously for 2 hours. The mixture was diluted with dichloromethane and water, and the organic layer was washed with water, dried and evaporated. Chromatography (silica, 5% methanol in dichloromethane) gave the title compound (0.67 g) as an off-white solid. $^1\text{H-NMR}$ (CDCl_3) δ 3.61 (1H, t), 4.82 (2H, d), 7.33 (1H, d), 7.45 (2H, m), 7.51 (2H, m), 7.85 (1H, dd), 7.78 (1H, d); MS (APCI+) found ($M+1$) = 220/222; $\text{C}_{12}\text{H}_{10}\text{ClNO}$ requires 219/221.

20 **Intermediate A13 — Ethyl 2-(4-chlorophenyl)-4-oxypyrimidine-5-carboxylate**




- Sodium ethoxide (11.12 ml, 2 equiv) as a 21% w/v solution in ethanol was added dropwise to a suspension of diethyl ethoxymalonate (3.03 ml, 1 equiv) and 4-chlorobenzamidine hydrochloride (4.23 g, 1 equiv) in ethanol (30 ml), then the mixture was heated to reflux for 4 hours. After cooling, the
- 25 solvent was removed in vacuo and the residue was triturated with ether. The solid was filtered off, then resuspended in water and acidified to pH 2. The product was filtered off, washed with water and dried; yield 2.94 g. $^1\text{H-NMR}$ (d_6 -DMSO) δ 1.29 (3H, t), 4.26 (2H, q), 7.65 (2H, m), 8.18 (2H, m), 8.65 (1H, s); MS (APCI-) found ($M-1$) = 277/279; $\text{C}_{13}\text{H}_{11}\text{ClN}_2\text{O}_3$ requires 278/280.


Intermediate A14 — Ethyl 2-(4-chlorophenyl)-4-chloropyrimidine-5-carboxylate



- 30 Oxalyl chloride (0.31 ml, 2 equiv) was added to Intermediate A13 (0.49 g) in dichloromethane (20 ml) with ice cooling, then the mixture was stirred for 3 hours with warming to room temperature. Evaporation of the volatile components gave the product as a white solid (2.94 g). $^1\text{H-NMR}$ (CDCl_3) δ 1.44 (3H, t), 4.48 (2H, q), 7.50 (2H, m), 8.45 (2H, m), 9.17 (1H, s); MS (APCI+) found ($M+1$) = 297; $\text{C}_{13}\text{H}_{10}\text{Cl}_2\text{N}_2\text{O}_2$ requires 296.
- 35

Intermediate A15 — Ethyl 2-(4-chlorophenyl)pyrimidine-5-carboxylate

















A mixture of Intermediate A14 (6.8 g, 1 equiv), zinc powder (1.79 g, 1.2 equiv), acetic acid (1.57 ml, 1.2 equiv) and THF (100 ml) was stirred at 60°C under argon for 18 hours, then a further portion of acetic acid (1 ml) and zinc (1.0 g) was added, and the reaction allowed to continue for a further 24 hours. The solvent was removed in vacuo, the residue was taken up in a mixture of dichloromethane and methanol, and undissolved zinc powder was removed by filtration. After evaporation of the solvent, the product crystallised from ethanol; yield 2.02 g. ¹H-NMR (CDCl₃) δ 1.44 (3H,t), 4.46 (2H,q), 7.48 (2H,m), 8.48 (2H,m), 9.30 (2H,s); MS (APCI+) found (M+1) = 263; C₁₃H₁₁ClN₂O₂ requires 262.

Intermediate A16 — 5-Hydroxymethyl-2-(4-trifluoromethylphenyl)pyrimidine

Intermediate A129 (0.96g) was hydrogenated over 10% palladium on charcoal (96 mg) in a mixture of triethylamine (2 ml) and ethanol (20 ml) for 90 mins at 1 atmosphere pressure. The catalyst was removed by filtration, the solvent was evaporated, and the residue was taken up in ethyl acetate and washed successively with aq. ammonium chloride and aq. sodium bicarbonate. Drying and evaporation gave the title compound (2.02 g). ¹H-NMR (CDCl₃) δ 4.82 (2H,s), 7.75 (2H,m), 8.57 (2H,m), 8.85 (2H,s); MS (APCI+) found (M+1) = 255; C₁₁H₉ClN₂O requires 254.

The following intermediates were made by the method of Intermediate A1:

No	Precursors	Structure	Name
A2 0	4-formylboronic acid, 4-fluoriodobenzene		4-(4-Fluorophenyl)benzaldehyde
A2 1	2-bromothiophene-5-carboxaldehyde, 4-fluorobenzeneboronic acid		5-(4-Fluorophenyl)-2-thiophene carboxaldehyde
A2 2	2-bromothiophene-5-carboxaldehyde, 4-chlorobenzeneboronic acid		5-(4-Chlorophenyl)-2-thiophene carboxaldehyde
A2 3	methyl 6-chloronicotinate, 4-fluorobenzeneboronic acid		Methyl 6-(4-fluorophenyl)-nicotinate
A2 4	methyl 6-chloronicotinate, 4-chlorobenzeneboronic acid		Methyl-6-(4-chlorophenyl)-nicotinate
A2 5	methyl 6-chloronicotinate, 3,4-dichlorobenzeneboronic acid		Methyl-6-(3,4-dichlorophenyl)-nicotinate

A2 6	4-bromobenzaldehyde, 4-trifluoromethylbenzene- boronic acid		4-(4-Trifluoromethylphenyl)- benzaldehyde
A2 7	4-bromo-2-fluorobenzaldehyde, 4-chlorobenzeneboronic acid		4-(4-Chlorophenyl)-2-fluoro- benzaldehyde
A2 8	4-formylbenzeneboronic acid, 4-chloro-2-fluoriodobenzene		4-(4-Chloro-2-fluorophenyl)- benzaldehyde
A2 9	4-formylbenzeneboronic acid, 4-chloro-3-fluoriodobenzene		4-(4-Chloro-3-fluorophenyl)- benzaldehyde
A3 0	4-methoxybenzeneboronic acid, 4-bromobenzaldehyde		4-(4-Methoxyphenyl)- benzaldehyde
A3 1	4-formylbenzeneboronic acid, 4-chloro-2-fluoriodobenzene		4-(2-Fluoro-4-chlorophenyl)- benzaldehyde
A3 2	4-formylbenzeneboronic acid, 2,4-bis(trifluoromethyl)- bromobenzene		4-(2,4-bis(trifluoromethyl)- phenyl)benzaldehyde
A3 3	4-formylbenzeneboronic acid, 4-bromo-3-fluorobenzo- trifluoride		4-(2-Fluoro-4-trifluoromethyl- phenyl)benzaldehyde
A3 4	4-bromo-2-fluorobenzaldehyde, 4-trifluoromethylbenzene- boronic acid		2-Fluoro-4-(4-trifluoromethyl- phenyl)benzaldehyde
A3 5	3-chloro-4-fluorobenzene- boronic acid, 4-bromobenzaldehyde		4-(3-Chloro-4-fluorophenyl)- benzaldehyde
A3 6	4-formylbenzeneboronic acid, 2,4-difluoriodobenzene		4-(2,4-Difluorophenyl)- benzaldehyde
A3 7	4-formylbenzeneboronic acid, 4-trifluoromethoxyiodobenzene		4-(4-Trifluoromethoxyphenyl)- benzaldehyde
A3 8	4-formylbenzeneboronic acid, 4-difluoromethoxyiodobenzene		4-(4-Difluoromethoxyphenyl)- benzaldehyde
A3 9	N-methyl-4-bromo-3- fluorobenzamide, 4-chlorobenzeneboronic acid		N-Methyl-4-(4-chlorophenyl)-3- fluorobenzamide
A4 0	methyl 6-chloronicotinate, 4- trifluoromethylbenzeneboronic acid		Methyl 6-(4-trifluoromethyl- phenyl)nicotinate
A4 1	2-methyl-5-bromopyridine, 4-chlorobenzeneboronic acid		2-methyl-5-(4-trifluoromethyl- phenyl)pyridine

A4 2	2-methyl-5-triflylpyridine, 4-trifluoromethylbenzeneboronic acid		2-methyl-5-(4-trifluoromethyl- phenyl)pyridine
A4 3	2-methyl-5-triflylpyridine, 4-trifluoromethoxybenzene- boronic acid		2-methyl-5-(4-trifluoromethoxy- phenyl)pyridine

The following intermediates were made by the method of Intermediate A2, using an appropriate amine in place of methylamine as necessary:

No	Precursor	Structure	Name
A4 4	Int. A20		N-Methyl-4-(4-fluorophenyl)benzylamine
A4 5	Int. A26		N-Methyl-4-(4-trifluoromethylphenyl)- benzylamine
A4 6	Int. A27		N-Methyl-4-(4-chlorophenyl)-2-fluoro- benzylamine
A4 7	Int. A28		N-Methyl-4-(4-chloro-2-fluorophenyl)- benzylamine
A4 8	Int. A1		N-Ethyl-4-(4-chlorophenyl)benzylamine
A4 9	Int. A131		5-Methylaminomethyl-2-(3,4-dichlorophenyl)- pyridine
A5 0	Int. A1		N-(Dimethylaminocarbonylmethyl)-4-(4- chlorophenyl)benzylamine
A5 1	Int. A1		N-(2-Methoxyethyl)-4-(4-chlorophenyl)- benzylamine
A5 2	Int. A26		N-(Dimethylaminocarbonylmethyl)-4-(4- trifluoromethylphenyl)benzylamine
A5 3	Int. A29		N-Methyl-4-(4-chloro-3-fluorophenyl)- benzylamine
A5 4	Int. A30		N-Methyl-4-(4-methoxyphenyl)benzylamine
A5 5	Int. A31		N-Methyl-4-(2-fluoro-4-chlorophenyl)- benzylamine
A5 6	Int. A32		N-Methyl-4-(2,4-bis(trifluoromethyl)phenyl)- benzylamine

A5 7	Int. A33		N-Methyl-4-(2-fluoro-4-trifluoromethylphenyl)-benzylamine
A5 8	Int. A34		N-Methyl-2-fluoro-4-(4-trifluoromethylphenyl)benzylamine
A5 9	Int. A35		N-Methyl-4-(3-chloro-4-fluorophenyl)-benzylamine
A6 0	Int. A36		N-Methyl-4-(2,4-difluorophenyl)benzylamine
A6 1	Int. A1		N-(2-Dimethylaminoethyl)-4-(4-chlorophenyl)-benzylamine
A6 2	Int. A1		N-(3-Dimethylaminopropyl)-4-(4-chlorophenyl)-benzylamine
A6 3	Int. A21		2-Methylaminomethyl-5-(4-fluorophenyl)-thiophene
A6 4	Int. A9		5-Methylaminomethyl-2-(4-fluorophenyl)-pyridine
A6 5	Int. A130		5-Methylaminomethyl-2-(4-chlorophenyl)-pyridine
A6 6	Int. A22		2-Methylaminomethyl-5-(4-chlorophenyl)-thiophene

A67: N-Methyl-4-phenylbenzylamine was made from commercially available 4-biphenylcarboxaldehyde.




The following intermediates were made by the method of Intermediate A3:

No.	Precursor	Structure	Name
A70	Int. A1		N-(2-Hydroxyethyl)-4-(4-chlorophenyl)-benzylamine
A71	Int. A26		N-(2-Hydroxyethyl)-4-(4-trifluoromethylphenyl)benzylamine
A72	Int. A26		N-(2-(dimethylamino)ethyl)-4-(4-trifluoromethylphenyl)benzylamine
A73	Int. A26		N-(2-(diethylamino)ethyl)-4-(4-trifluoromethylphenyl)benzylamine
A74	Int. A1		N-(2-(diisopropylamino)ethyl)-4-(4-chlorophenyl)benzylamine





A75	Int. A1		N-(2-(1-pyrrolidino)ethyl)-4-(4-chlorophenyl)benzylamine
A76	Int. A1		N-(2-(4-morpholino)ethyl)-4-(4-chlorophenyl)benzylamine
A77	Int. A1		N-(2-(4-methyl-1-piperazino)ethyl)-4-(4-chlorophenyl)benzylamine
A78	Int. A26		N-(3-(dimethylamino)propyl)-4-(4-(trifluoromethyl)phenyl)benzylamine
A79	Int. A1		N-(3-(diethylamino)propyl)-4-(4-chlorophenyl)benzylamine
A80	Int. A1		N-(3-(1-pyrrolidino)propyl)-4-(4-chlorophenyl)benzylamine
A81	Int. A1		N-(3-(1-piperidino)propyl)-4-(4-chlorophenyl)benzylamine
A82	Int. A1		N-(3-(4-morpholino)propyl)-4-(4-chlorophenyl)benzylamine
A83	Int. A1		N-(3-(4-methyl-1-piperazino)propyl)-4-(4-chlorophenyl)benzylamine
A84	Int. A1		N-(2-(t-butoxycarbonylamino)ethyl)-4-(4-chlorophenyl)benzylamine
A85	Int. A1		N-(N'-(2-(t-butoxycarbonyl)-N'-methylamino)ethyl)-4-(4-chlorophenyl)benzylamine
A86	Int. A1		N-(N'-(2-(t-butoxycarbonyl)-N'-ethylamino)ethyl)-4-(4-chlorophenyl)benzylamine
A87	Int. A26		N-(N'-(2-(t-butoxycarbonyl)-N'-ethylamino)ethyl)-4-(4-(trifluoromethyl)phenyl)benzylamine
A88	Int. A1		N-(3-(t-butoxycarbonylamino)propyl)-4-(4-chlorophenyl)benzylamine
A89	Int. A1		N-(N'-(3-(t-butoxycarbonyl)-N'-methylamino)propyl)-4-(4-(trifluoromethyl)phenyl)benzylamine
A90	Int. A1		N-(2-(4-t-butoxycarbonyl-1-piperazino)ethyl)-4-(4-chlorophenyl)benzylamine
A91	Int. A1		N-(2-(2-oxo-4-t-butoxycarbonyl-1-piperazino)ethyl)-4-(4-chlorophenyl)benzylamine

A92	Int. A37		N-(2-(diethylamino)ethyl)-4-(4-(trifluoromethoxyphenyl)benzylamine
A93	Int. A38		N-(2-(diethylamino)ethyl)-4-(4-(difluoromethoxyphenyl)benzylamine
A94	Int. A1		N-(2-(N'-(2-hydroxyethyl)-N'-ethylamino)ethyl)-4-(4-chlorophenyl)benzylamine
A95	Int. A1		N-(2-(bis-(2-hydroxyethyl)amino)ethyl)-4-(4-chlorophenyl)benzylamine
A96	Int. A26		N-((4-morpholino)carbonylmethyl)-4-(4-(trifluoromethyl)phenyl)benzylamine
A97	Int. A26		N-((4-methyl-1-piperazino)carbonylmethyl)-4-(4-(trifluoromethyl)phenyl)benzylamine
A98	Int. A133		N-(2-(diethylamino)ethyl)-5-(4-chlorophenyl)pyrid-2-ylmethylamine
A99	Int. A134		N-(2-(diethylamino)ethyl)-5-(4-(trifluoromethyl)phenyl)pyrid-2-ylmethylamine
A100	Int. A132		N-(2-(dimethylamino)ethyl)-2-(4-(trifluoromethyl)phenyl)pyrid-5-ylmethylamine
A101	Int. A130		N-(2-(diethylamino)ethyl)-2-(4-chlorophenyl)pyrid-5-ylmethylamine
A102	Int. A135		N-methyl-5-(4-(trifluoromethoxy)phenyl)pyrid-2-ylmethylamine
A103	Int. A130		N-(2-hydroxyethyl)-2-(4-chlorophenyl)pyrid-5-ylmethylamine
A104	Int. A136		N-(2-(diethylamino)ethyl)-2-(4-chlorophenyl)pyrimid-5-ylmethylamine
A105	Int. A137		N-(2-(diethylamino)ethyl)-2-(4-(trifluoromethyl)phenyl)pyrimid-5-ylmethylamine
A106	Int. A132		N-(2-(diethylamino)ethyl)-2-(4-(trifluoromethyl)phenyl)pyrid-5-ylmethylamine
A107	Int. A1		N-(2-(1-piperidino)ethyl)-4-(4-chlorophenyl)benzylamine
A108	Int. A26		N-(2-(1-piperidino)ethyl)-4-(4-(trifluoromethyl)phenyl)benzylamine





The following intermediates were made by the method of Intermediate A4:

No.	Precursor	Structure	Name
A110	Int. A20		4-(4-Fluorophenyl)benzyl alcohol
A111	Int. A26		4-(4-Trifluoromethylphenyl)benzyl alcohol
A112	Int. A22		5-Hydroxymethyl-2-(4-chlorophenyl)thiophene

The following intermediates were made by the method of Intermediate A5:

No.	Precursor	Structure	Name
A11 5	Int. A110		N-(4-(4-Fluorophenyl)benzyl)phthalimide
A11 6	Int. A111		N-(4-(4-Trifluoromethylphenyl)benzyl)-phthalimide
A11 7	Int. A8		N-(2-(4-Fluorophenyl)pyrid-5-ylmethyl)-phthalimide
A11 8	Int. A112		N-(5-(4-Chlorophenyl)thien-2-ylmethyl)-phthalimide

The following intermediates were made by the method of Intermediate A6:

No.	Precursor	Structure	Name
A12 0	Int. A115		4-(4-Fluorophenyl)benzylamine
A12 1	Int. A116		4-(4-Trifluoromethylphenyl)benzylamine
A12 2	Int. A117		5-Aminomethyl-2-(4-fluorophenyl)pyridine
A12 3	Int. A118		5-Aminomethyl-2-(4-chlorophenyl)thiophene

A124: 4-Phenylbenzylamine is commercially available.

The following intermediates were made by the method of Intermediate A8:

No.	Precursor	Structure	Name
A125	Int. A24		5-Hydroxymethyl-2-(4-chlorophenyl)pyridine
A126	Int. A25		5-Hydroxymethyl-2-(3,4-dichlorophenyl)pyridine
A127	Int. A40		5-Hydroxymethyl-2-(4-trifluoromethylphenyl)pyridine
A128	Int. A15		5-Hydroxymethyl-2-(4-chlorophenyl)pyrimidine
A129	Int. A150		4-chloro-5-hydroxymethyl-2-(4-trifluoromethylphenyl)pyrimidine


The following intermediates were made by the method of Intermediate A9:

No.	Precursor	Structure	Name
A130	Int. A125		5-Formyl-2-(4-chlorophenyl)pyridine
A131	Int. A126		5-Formyl-2-(3,4-dichlorophenyl)pyridine
A132	Int. A127		5-Formyl-2-(4-trifluoromethylphenyl)pyridine
A133	Int. A12		2-Formyl-5-(4-chlorophenyl)pyridine
A134	Int. A140		2-Formyl-5-(4-trifluoromethylphenyl)pyridine
A135	Int. A141		2-Formyl-5-(4-trifluoromethoxyphenyl)pyridine
A136	Int. A128		5-Formyl-2-(4-chlorophenyl)pyrimidine
A137	Int. A16		5-Formyl-2-(4-trifluoromethylphenyl)pyrimidine


The following intermediates were made by the method of Intermediate A12:

No.	Precursor	Structure	Name
A140	Int. A134		2-hydroxymethyl-5-(4-trifluoromethylphenyl)pyridine
A141	Int. A135		2-hydroxymethyl-5-(4-trifluoromethoxyphenyl)pyridine


The following intermediate was made by the method of Intermediate A13:

No.	Precursors	Structure	Name
A14 5	diethyl ethoxymalonate, 4-trifluoromethyl- benzamidinium.HCl		Ethyl 2-(4-trifluoromethylphenyl)-4-oxopyrimidine-5-carboxylate


The following intermediate was made by the method of Intermediate A14:


No.	Precursor	Structure	Name
A15 0	Int. A145		Ethyl 2-(4-trifluoromethylphenyl)-4-chloropyrimidine-5-carboxylate

The following intermediate was made by the method of Intermediate A7:


No.	Precursor	Structure	Name
A15 5	Int. A45		N-Methyl-N-(4-(4-trifluoromethylphenyl)benzyl)bromoacetamide

The following intermediate was made by the method of Intermediate A11:


No.	Precursor	Structure	Name
A16 0	Int. A26		N-(Ethoxycarbonylmethyl)-4-(4-trifluoromethylphenyl)benzylamine

Intermediate B1 — Ethyl 2-trifluoromethyl-4-oxypyrimidine-5-carboxylate


- To a solution of trifluoroacetamidine (36.5 g) and diethyl ethoxymalonate (71 g) in ethanol (300 ml) was added a solution of sodium ethoxide in ethanol (21 wt %, 109.5 ml) over 5 min. The mixture was heated at reflux for 6 h, then cooled, concentrated and water (200 ml) added. The resulting solid was collected by filtration, washed with cold water (50 ml) and diethyl ether (2 x 100 ml) and then suspended in water (400 ml). Dichloromethane (300 ml) was added and the mixture acidified with dilute HCl (2.5 M, 125 ml). The organic extract was washed with water, dried (MgSO₄) and evaporated to give the title compound as a buff solid (53 g, 68%). ¹H-NMR (CDCl₃) δ 12.00 (1H, br s), 9.18 (1H, s), 4.57 (2H, q), 1.49 (3H, t); ¹³C-NMR (CDCl₃) δ 171.4, 167.3, 160.4, 160.2 (q, J = 38 Hz), 118.8 (q, J = 276 Hz), 109.2, 63.8, 14.0; MS (APCI+) found (M+1) = 237, C₈H₇F₃N₂O₃ requires 236.

Intermediate B2 — Ethyl 2-trifluoromethyl-4-chloropyrimidine-5-carboxylate

- To a solution of ethyl 4-hydroxy-2-trifluoromethylpyrimidine-5-carboxylate (51.8 g) in dichloromethane (600 ml) cooled in an ice bath was added oxalyl chloride (57.4 ml) followed by dimethylformamide (0.2 ml). The mixture was stirred at room temperature for 16 h and then evaporated. Toluene was added and evaporated. The residue was dissolved in dichloromethane, washed with water, dried (MgSO₄) and evaporated to give the title compound as an orange oil (55.7 g, 100%). ¹H-NMR (CDCl₃) δ 9.25 (1H, s), 4.51 (2H, q), 1.46 (3H, t); ¹³C-NMR (CDCl₃) δ 162.0 (2C), 161.1, 158.1 (q, J = 39 Hz), 127.0, 118.9 (q, J = 276 Hz), 63.5, 14.4.


Intermediate B3 — Ethyl 2-trifluoromethylpyrimidine-5-carboxylate

- A mixture of ethyl 4-chloro-2-trifluoromethylpyrimidine-5-carboxylate (55.7 g), 10% palladium on carbon (0.3 g), ethanol (1000 ml) and *N,N*-diisopropylethylamine (90 ml) was shaken under hydrogen pressure maintained at 1 atmosphere for 2 h. The catalyst was then filtered off and the solvents evaporated. The residue was dissolved in dichloromethane, washed with ammonium chloride solution, then water, dried (MgSO₄) and evaporated to give the title compound as a buff solid (48 g, 100%). ¹H-NMR (CDCl₃) δ 9.42 (2H, s), 4.51 (2H, q), 1.45 (3H, t); ¹³C-NMR (CDCl₃) δ 162.7, 159.4 (2C), 159.3 (q, J = 37 Hz), 126.3, 119.6 (q, J = 275 Hz), 62.9, 14.4.

Intermediate B4 — 2-Trifluoromethyl-5-formylpyrimidine


To a solution of ethyl 2-trifluoromethylpyrimidine-5-carboxylate (5.17 g) in toluene (120 ml) cooled in dry ice/acetone was added a solution of diisobutylaluminium hydride (25 wt %, 31 ml) over 15 min.

- 5 The mixture was stirred at -78°C for 45 min, then dilute HCl (2M, 120 ml) was added cautiously. After allowing the mixture to warm to room temperature diethyl ether was added. The organic phase was separated, washed with water, then brine, dried (MgSO₄) and evaporated to give the title compound as a colourless solid (3.46 g, 84%). ¹H-NMR (CDCl₃) δ 10.29 (1H, s), 9.37 (2H, s); ¹³C-NMR (CDCl₃) δ 187.7, 159.6 (q, J = 38 Hz), 159.1 (2C) 129.6, 119.2 (q, J = 276 Hz).


Intermediate B5 – 1-(2-Methoxyethyl)pyrazole

2-Bromomethyl methyl ether (10.22 g) was added dropwise with stirring to a mixture of pyrazole (5.0 g), finely powdered potassium hydroxide (8.25 g) and tetrabutylammonium bromide (1.19 g) with occasional ice cooling to keep the temperature below 10°C. The mixture was allowed to stand at room temperature for 48 hours, then columned on silica and eluted with ether. Product fractions were evaporated to a pale green oil (7.27 g). ¹H-NMR (CDCl₃) δ 3.33 (3H,s), 3.75 (2H,t), 4.30 (2H,t), 6.25 (1H,s), 7.47 (1H,d), 7.52 (1H,d); MS (APCI+) found (M+H) = 127; C₆H₁₀N₂O requires 126.


15

Intermediate B6 – 1-(2-Methoxyethyl)pyrazole-4-carboxaldehyde


- 20 A solution of 1-(2-methoxyethyl)pyrazole (7.27 g) in dry DMF (11.4 ml) was heated to 90°C, then phosphorus oxychloride (5.4 ml) was added dropwise over 1 hour, maintaining the temperature between 95-100°C. After heating for a further 2 hours, the mixture was cooled and poured onto ice. Sodium hydroxide was added to adjust the mixture to pH 4, then the product was extracted into dichloromethane. Drying and evaporation of the organic extracts yielded a brown oil (7.19 g). ¹H-NMR (CDCl₃) δ 3.34 (3H,s), 3.75 (2H,m), 4.32 (2H,m), 7.98 (1H,s), 8.02 (1H,s), 9.85 (1H,s).
- 25

Intermediate B10 – 3-(1-Methylpyrazol-4-yl)acrylic acid


- A mixture of 1-methylpyrazole-4-carboxaldehyde (85.55 g), malonic acid (80.85 g), pyridine (69.2 ml) and piperidine (1.5 ml) was heated to 110°C under argon for 4 hours. After cooling, water (100 ml) was added, followed by aqueous ammonia (75 ml) to obtain a clear solution, which was acidified to pH1 with hydrochloric acid. The resulting solid was filtered off, washed with water and dried to obtain the title compound (93.5 g). ¹H-NMR (d₆-DMSO) δ 3.83 (3H,s), 6.18 (1H,d), 7.44 (1H,d), 7.83 (1H,s), 8.07 (1H,s). (APCI) found (M+H)=153. C₇H₈N₂O₂ requires 152.
- 30

Intermediate B11 – Methyl 3-(1-methylpyrazol-4-yl)acrylate

3-(1-Methylpyrazol-4-yl)acrylic acid (86.8 g) was added to a solution of sulphuric acid (20 ml) in methanol (690 ml), and the mixture refluxed for 2.5 hours, cooled, then poured onto ice. The acid was neutralised with aqueous sodium hydroxide and the product extracted into dichloromethane, which was dried and evaporated. Crystallisation from ether/petrol gave methyl 3-(1-methylpyrazol-4-yl)acrylate (89.0 g). ¹H-NMR (d₆-DMSO) δ 3.77 (3H,s), 3.91 (3H,s), 6.16 (1H,d), 7.54 (1H,s), 7.56 (1H,d), 7.69 (1H,s). (APCI) found (M+H)=167. C₈H₁₀N₂O₂ requires 166.

Intermediate B12 – Ethyl 3-(5-pyrimidinyl)acrylate


A mixture of 5-bromopyrimidine (5.93g), ethyl acrylate (5.08g), palladium acetate (0.112g), triphenyl phosphine (0.23g) and triethylamine (4.5g) was stirred at 150°C in a pressure vessel for 6 hours. After cooling overnight, water (50ml) was added to the dark residue, and the product was extracted into toluene. Drying, charcoaling and evaporation gave a pale oil, which was triturated with pet. ether to obtain ethyl 3-(5-pyrimidinyl)acrylate (4.78g). ¹H-NMR (CDCl₃) δ 1.36 (3H,t), 4.27 (2H,q), 6.59 (1H,d), 7.62 (1H,d), 8.88 (2H,s), 9.20 (1H,s).

Intermediate B13 – Ethyl 3-(2-methoxypyrimidin-5-yl)acrylate



A mixture of 2-methoxy-5-bromopyrimidine (75.43 g, 0.399 mol), ethyl acrylate (47.5 ml, 0.439 mol), palladium (II) acetate (1.07 g, 0.0048 mol), tri-*o*-tolylphosphine (2.92 g, 0.0096 mol) and triethylamine (84 ml) were heated at 135°C (oil bath temperature) with stirring under argon for 12 h. After allowing to cool the solid mass was dissolved in water and ethyl acetate, filtered, and the aqueous phase separated and further extracted with ethyl acetate. The combined extracts were washed with saturated aqueous ammonium chloride, dried (MgSO₄) and evaporated. The solid thus obtained was triturated with ether/light petrol (1:3, 350 ml), filtered, washed and dried, yield 52.41 g (63%). ¹H-NMR (CDCl₃) δ 1.33 (3H, t), 4.06 (3H, s), 4.28 (2H, q), 6.45 (1H, d), 7.58 (1H, d), 8.67 (2H, s); MS (APCI+) found (M+H) = 209; C₁₀H₁₂N₂O₃ requires 208.

The following intermediates were prepared by the method of intermediate B10 (Knoevenagel):



No	Precursors	Structure	Name
B14	Int. B4		3-(2-Trifluoromethylpyrimidin-5-yl)acrylic acid

B1 5	Int. B6		3-(1-(2-Methoxyethyl)pyrazol-4-yl)acrylic acid
-----------------------	---------	---	--


The following intermediates were prepared by the method of intermediate B11 (esterification):

No	Precursors	Structure	Name
B1 6	Int. B14		Methyl 3-(2-trifluoromethylpyrimidin-5-yl)acrylate
B1 7	Int. B15		Methyl 3-(1-(2-methoxyethyl)pyrazol-4-yl)acrylate

The following intermediates were prepared by the method of intermediate B13 (acrylate Heck):


No	Precursors	Structure	Name
B1 8	2-dimethylamino-5-bromopyrimidine		Ethyl 3-(2-dimethylamino-5-pyrimidyl)-acrylate
B1 9	2-(4-morpholino)-5-bromopyrimidine		Ethyl 3-[2-(4-morpholino)-5-pyrimidyl]-acrylate

Intermediate B20 – Methyl 3-(1-methylpyrazol-4-yl)propionate



- 5 A solution of methyl 3-(1-methylpyrazol-4-yl)acrylate (181 g) in methanol (2 litres) was hydrogenated over 10% palladium on charcoal (5.2 g) at 50°C/50psi until uptake ceased. The catalyst was filtered off, the methanol was removed in vacuo, finally by azeotrope with toluene. The title compound was obtained as an oil (179 g). ¹H-NMR (d₆-DMSO) δ 2.56 (2H,t), 2.79 (2H,t), 3.67 (3H,s), 3.85 (3H,s), 7.17 (1H,s), 7.31 (3H,s). (APCI) M+H=169. C₈H₁₂N₂O₂ requires 168.

10 Intermediate B21 – Ethyl 3-(2-methoxypyrimidin-5-yl)propanoate



- 15 A suspension of ethyl 3-(2-methoxypyrimidin-5-yl)acrylate (52.4 g, 0.252 mol) in ethanol (400 ml) and triethylamine (50 ml) was treated with 10% palladium on carbon (3 g) and hydrogenated at 50 psi for 1.75 h. The catalyst was filtered off through hyflo and the filtrate evaporated. The residue was dissolved in dichloromethane, washed twice with saturated aqueous ammonium chloride, dried

(MgSO₄) and evaporated to an oil, yield 41.2 g (78%). ¹H-NMR (CDCl₃) δ 1.23 (3H, t), 2.61 (2H, t), 2.87 (2H, t), 3.99 (3H, s), 4.13 (2H, q), 8.39 (2H, s); MS (APCI+) found (M+H) = 211; C₁₀H₁₄N₂O₃ requires 210.


- 5 The following intermediates were made by the method of Intermediate B20 (ethyl esters in ethanol solvent, methyl esters in methanol):

No	Precursors	Structure	Name
B2 2	Int. B12		Ethyl 3-(5-pyrimidyl)propionate
B2 3	Int. B17		Methyl 3-(1-(2-methoxyethyl)pyrazol-4-yl)propionate

The following intermediates were made by the method of Intermediate B21 (ethyl esters in ethanol solvent, methyl esters in methanol):

No	Precursors	Structure	Name
B2 4	Int. B16		Methyl 3-(2-trifluoromethylpyrimidin-5-yl)propanoate
B2 5	Int. B18		Ethyl 3-(2-dimethylamino-5-pyrimidyl)propionate
B2 6	Int. B19		Ethyl 3-[2-(4-morpholino)-5-pyrimidyl]propionate


Intermediate B30 – 5-((1-Methylpyrazol-4-yl)methyl)-2-thiouracil



- 10 A solution of methyl 3-(1-methylpyrazol-4-yl)propionate (170 g) and methyl formate (131 ml) in dry diethyl ether (2000 ml) was added dropwise over 2 hours to a stirred, ice-cooled solution of potassium tert-butoxide (284g) in dry THF (1800 ml) under argon. The mixture was then allowed to warm to room temperature and stirring continued for 16 hours. The solvents were evaporated in vacuo to a pale solid. Methanol (2500 ml) and thiourea (154 g) were added, and the mixture was heated to 50°C for 16
- 15 hours with vigorous stirring. The solvent was evaporated, and the pale brown solid residue was taken up in water (750 ml). The ice-cooled solution was adjusted to pH3 with hydrochloric acid, stirred for 2 hours in the ice bath, then the precipitate was filtered off and washed with water and ether to obtain the


title compound (120 g). $^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$) δ 3.33 (3H,s), 3.75 (3H,s), 7.15 (1H,s), 7.23 (1H,s), 7.46 (1H,s), 12.2 (1H,br s), 12.4 (1H,br s). (APCI) $\text{M}+\text{H}=223$. $\text{C}_9\text{H}_{10}\text{N}_4\text{OS}$ requires 222.

Intermediate B31 – Ethyl 2-formyl-3-(5-pyrimidyl)propionate



- 5 A mixture of ethyl 3-(5-pyrimidyl)propionate (2.28g) and ethyl formate (1.41ml) dissolved in dry dimethoxyethane (5ml) was added dropwise over 30 min to a suspension of sodium hydride (60%, 4.0g) in DME (5 ml) under nitrogen, keeping the temperature below 0°C . Stirring was continued for a further 24h, then the mixture was poured onto ice and washed with ether. The aqueous layer was adjusted to pH 7, then evaporated and the residue extracted with acetone. Filtration and evaporation
- 10 gave crude product, which was taken up in ethyl acetate, charcoaled, dried and evaporated to give ethyl 2-formyl-3-(5-pyrimidyl)propionate. Like other compounds of this type, this proved difficult to characterise and was used without further purification.





Intermediate B32 – 5-(Pyrimid-5-ylmethyl)-2-thiouracil




- 15 Sodium (0.25 g) was dissolved in ethanol (5 ml), thiourea (0.77 g) added, and the mixture stirred under reflux for 1 hour. A solution of ethyl 2-formyl-3-(5-pyrimidyl)propionate (1.99 g) in ethanol (5 ml) was added slowly, and reflux continued for 18 hours. The solvent was evaporated, and the residue taken up in water and washed with dichloromethane. The aqueous solution was acidified to pH 5, and the precipitate filtered off, washed with water and dried to obtain 5-(pyrimid-5-ylmethyl)-2-thiouracil
- 20 (0.71 g). $^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$) δ 3.58 (2H,s), 7.54 (1H,s), 8.70 (2H,s) and 9.02 (1H,s). MPt $265-6^\circ\text{C}$

The following intermediates were made by the method of Intermediate B30 (thiouracils):

No	Precursors	Structure	Name
B3 3	Int. B21		5-(2-Methoxypyrimidin-5-ylmethyl)-2-thiouracil
B3 4	Int. B24		5-(2-Trifluoromethylpyrimidin-5-ylmethyl)-2-thiouracil
B3 5	Int. B26		5-(2-(4-Morpholino)pyrimidin-5-ylmethyl)-2-thiouracil


B3 6	Int. B25		5-(2-Dimethylaminopyrimid-5-ylmethyl)-2-thiouracil
B3 7	Int. B23		5-((1-(2-Methoxyethyl)pyrazol-4-yl)methyl)-2-thiouracil
B3 8	ethyl 3-phenyl-propionate		5-Benzyl-2-thiouracil
B3 9	ethyl 3-(4-chlorophenyl)-propionate		5-(4-Chlorobenzyl)-2-thiouracil

Intermediate B40 – 2-(Methoxymethylene)-3-(2-methoxypyrimidin-5-yl)propionic acid, mixed methyl/ethyl esters




- To a stirring suspension of sodium hydride (0.83 g of a 60% dispersion in oil) in anhydrous 1, 2-dimethoxyethane (6 ml) was added dropwise a solution of methyl formate (1.54 ml) and ethyl 3-(2-methoxypyrimid-5-yl)propionate (3.5 g) in anhydrous 1,2-dimethoxyethane (6 ml) at such a rate as to maintain the reaction temperature at 25-30°C. After 1 h, ether was added and the precipitated oil allowed to settle. The solution was decanted off and replaced with fresh ether, and the oil slowly solidified. The solid 2-(hydroxymethylene) derivative was filtered, washed with ether and dried, yield 3.8 g. A 1.33 g portion was suspended in dimethyl formamide (10 ml) together with anhydrous potassium carbonate (1.15 g), and a solution of dimethyl sulphate (0.48 ml) in dimethylformamide (10 ml) was added dropwise with stirring over 30 min. After 16 h the solvent was evaporated and the residue treated with water and extracted with ethyl acetate. The extracts were washed with water, dried (MgSO₄) and evaporated to give the product as an oil, yield 0.91 g. ¹H-NMR (CDCl₃) δ 1.23 (3H, t), 3.46 (2H, s), 3.69 (3H, s, methyl ester), 3.88 (3H, s), 3.97 (3H, s), 4.16 (2H, q), 7.39 (1H, s), 8.40 (2H, s). 3:2 ratio of methyl:ethyl esters. MS (APCI+) found (M+1) = 253, 239 (ethyl and methyl esters); C₁₂H₁₆N₂O₄ requires 252, C₁₁H₁₄N₂O₄ requires 238.

Intermediate B41 – 2-(Methoxymethylene)-3-(2-methoxypyrimidin-5-yl)propionic acid



- A suspension of the mixed esters of Intermediate B40 (0.9 g) in 2M aqueous sodium hydroxide (3.6 ml) was stirred at ambient temperature for 16 h to give a clear solution. This was diluted with water, extracted with dichloromethane and evaporated to about half volume, then acidified to pH 3-4 (2M hydrochloric acid) when the product crystallised out. The white solid was filtered, washed with ice-cold water and dried, yield 0.46 g. ¹H-NMR (CDCl₃) δ 3.43 (2H, s), 3.91 (3H, s), 3.99 (3H, s), 7.49 (1H, s), 8.42 (2H, s); MS (APCI+) found (M+1) = 225, C₁₀H₁₂N₂O₄ requires 224.

Intermediate B42 – 1-(3-Ethoxycarbonylprop-1-yl)-5-(2-methoxypyrimidin-5-ylmethyl)-2-thiouracil




- Oxalyl chloride (3.94ml, 2 equiv) was added over 5min to a solution of Intermediate B41 (5.0g, 1 equiv) and DMF (1 drop) in dry dichloromethane (50ml), then the mixture was stirred under argon for 4h. The solvent was evaporated, and the residue twice taken up in dichloromethane and re-evaporated to remove volatile impurities. The acid chloride was dissolved in acetonitrile (100ml) and treated with dry, powdered potassium thiocyanate (3.2g, 1.5 equiv). The resulting suspension was stirred under argon for 24h, then evaporated to dryness. The residue was suspended in dry DMF (100ml), cooled to 10°C, ethyl 4-aminobutyrate hydrochloride (4.60g, 1.25 equiv) and triethylamine (7.34ml, 2.4 equiv) added, and the mixture stirred at room temperature for 20h. Sodium methoxide solution (prepared by dissolving sodium (1.26g, 2.5 equiv) in methanol (25ml)) was added to the DMF solution, and the mixture heated to 110°C for 2h. The solvent was evaporated, water was added, and acidified to pH 5 with acetic acid, then the product extracted into ethyl acetate. Drying, evaporation, and trituration with ether gave a pale brown solid (4.55g). Some ether/ester exchange took place during the reaction, and the title compound was obtained mixed with ca 30% of the corresponding 2-ethoxypyrimidine. ¹H-NMR (DMSO-d₆) δ 1.17 (3H, t), 1.31 (t, ethoxy), 1.97 (2H, t), 2.36 (2H, t), 3.49 (s, ethoxy), 3.51 (s, methoxy), 3.88 (s, methoxy), 4.04 (2H, q), 4.16 (2H, t), 4.31 (q, ethoxy), 7.81 (1H, s), 8.48 (s, ethoxy), 8.50 (s, methoxy), 12.50 (bs, NH) (light brown solid).

- 20 The following intermediates were prepared by the method of intermediate B40:



No.	Precursors	Structure	Name
B43	Int. B20		Methyl 2-(methoxymethylene)-3-(1-methylpyrazol-4-yl)propionate
B44	Int. B26		Ethyl 2-(methoxymethylene)-3-(2-(4-morpholino)pyrimidin-5-yl)propionate

The following intermediates were prepared by the method of intermediate B41:


No.	Precursor	Structure	Name
B45	Int. B43		2-(Methoxymethylene)-3-(1-methylpyrazol-4-yl)propionic acid

B4 6	Int. B44		2-(Methoxymethylene)-3-(2-(4-morpholino)pyrimidin-5-yl)propionic acid
-----------------	-----------------	---	--

The following intermediates were prepared by the method of intermediate B42:


No	Precursor s	Structure	Name
B4 7	Int. B45		1-(3-Ethoxycarbonylprop-1-yl)-5-(1-methylpyrazol-4-ylmethyl)-2-thiouracil
B4 8	Int. B46		1-(3-Ethoxycarbonylprop-1-yl)-5-(2-(4-morpholino)pyrimidin-5-ylmethyl)-2-thiouracil

Intermediate B50 – 2-(4-Fluorobenzylthio)-5-((1-methylpyrazol-4-yl)methyl)pyrimidin-4-one



- A mixture of 5-((1-methylpyrazol-4-yl)methyl)-2-thiouracil (118 g), 4-fluorobenzyl chloride (76.8 g), potassium carbonate (183.5 g), and dry DMF (100 ml) was stirred under argon at 80°C for 16 hours, then cooled and evaporated. The solid residue was suspended in water (1500 ml) with vigorous stirring, then acidified to pH2 with hydrochloric acid and stirred for a further hour. The white solid was filtered off and washed with water and ether to obtain the title compound (168 g). ¹H-NMR (d6-DMSO) δ 3.47 (2H, s), 3.81 (3H, s), 4.41 (2H, s), 7.19 (2H, s), 7.29 (1H, s), 7.48 (3H, m), 7.84 (1H, s), 12.74 (1H, br.s); MS (APCI+) found (M+1) = 331; C₁₆H₁₅FN₄OS requires 330.

Intermediate B51 – 1-(3-Ethoxycarbonylprop-1-yl)-2-(4-fluorobenzylthio)-5-(2-methoxy-pyrimidin-5-ylmethyl)pyrimidin-4-one



- Intermediate B24 (1.0g, 1 equiv) in dichloromethane (60ml) was treated with diisopropylethylamine (0.63ml, 1.3 equiv) followed by 4-fluorobenzyl bromide (0.38ml, 1.1 equiv), giving an orange solution which was stirred under argon for 4h, then washed with water, dried and evaporated. Chromatography (silica, 5% ethanol in ethyl acetate) eluted first the 2-ethoxypyrimidine impurity, followed by the title compound (0.35g). ¹H-NMR (CDCl₃) δ 1.25 (3H, t), 2.03 (2H, m), 2.33 (2H, t), 3.64 (2H, s), 3.83 (2H, m), 3.99 (3H, s), 4.12 (2H, q), 4.47 (2H, s), 6.98 (3H, m), 7.37 (2H, m), 8.46 (2H, s); MS (APCI+) M+1=473; C₂₃H₂₅FN₄O₄S requires 472 (pale waxy solid).

The following intermediates were prepared by the method of intermediate B50:


No	Precursors	Structure	Name
B5 2	Int. B32		2-(4-Fluorobenzylthio)-5-((pyrimidin-5-yl)methyl)pyrimidin-4-one
B5 3	Int. B33		2-(4-Fluorobenzylthio)-5-((2-methoxypyrimidin-5-yl)methyl)pyrimidin-4-one
B5 4	Int. B34		2-(4-Fluorobenzylthio)-5-((2-(trifluoromethyl)pyrimidin-5-yl)methyl)pyrimidin-4-one
B5 5	Int. B35		2-(4-Fluorobenzylthio)-5-((2-(4-morpholinomethyl)pyrimidin-5-yl)methyl)pyrimidin-4-one
B5 6	Int. B36		2-(4-Fluorobenzylthio)-5-((2-(dimethylaminomethyl)pyrimidin-5-yl)methyl)pyrimidin-4-one
B5 7	Int. B33 + 3,4-difluorobenzyl chloride		2-(3,4-Difluorobenzylthio)-5-((2-methoxypyrimidin-5-yl)methyl)pyrimidin-4-one
B5 8	Int. B37		2-(4-Fluorobenzylthio)-5-((1-(2-methoxyethyl)pyrazol-4-yl)methyl)pyrimidin-4-one
B5 9	Int. B30 + benzyl chloride		2-Benzylthio-5-(1-methylpyrazol-4-yl)methyl)pyrimidin-4-one
B6 0	Int. B30 + 2,6-dimethyl-4- chloromethyl- pyridine		2-(2,6-Dimethylpyridin-4-yl)methylthio-5-(1-methylpyrazol-4-yl)methyl)pyrimidin-4-one
B6 1	Int. B33 + 2,6-dichloro-4- chloromethyl- pyridine		2-(2,6-Dichloropyridin-4-yl)methylthio-5-(2-methoxypyrimidin-5-yl)methyl)pyrimidin-4-one

B6 2	Int. B33 + 3-chloro- methylpyridine		2-(Pyrid-3-yl)methylthio-5-(2-methoxy- pyrimid-5-ylmethyl)pyrimidin-4-one
B6 3	Int. B33 + 4-chloro- methyl-2- methylthiazole		2-(2-Methylthiazol-4-yl)methylthio-5-(2- methoxypyrimid-5-ylmethyl)pyrimidin-4- one
B6 4	Int. B38		2-(4-Fluorobenzylthio)-5-benzylpyrimidin- 4-one
B6 5	Int. B39		2-(4-Fluorobenzylthio)-5-(4-chlorobenzyl)- pyrimidin-4-one
B6 6	Int. B30 + 3,4-difluoro- benzyl chloride		2-Benzylthio-5-(1-methylpyrazol-4-yl- methyl)pyrimidin-4-one

The following intermediates were prepared by the method of intermediate B51:

No	Precursors	Structure	Name
B6 7	Int. B47		1-(3-Ethoxycarbonylprop-1-yl)-2-(4-fluoro- benzyl)thio-5-(1-methylpyrazol-4-ylmethyl)- pyrimidin-4-one
B6 8	Int. B48		1-(3-Ethoxycarbonylprop-1-yl)-2-(4-fluoro- benzyl)thio-5-(2-(4-morpholino)pyrimid-5-yl- methyl)pyrimidin-4-one


Intermediate B70 – 1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-((1-methylpyrazol-4-yl)methyl)pyrimidin-4-one



- 5 A mixture of 2-(4-fluorobenzylthio)-5-((1-methylpyrazol-4-yl)methyl)pyrimidin-4-one (175 g), tert-butyl iodoacetate (128.3 g), diisopropylethylamine (101.5 ml) and dichloromethane (200 ml) was stirred at room temperature under argon for 48 hours. The solution was washed with aq. sodium bicarbonate then with aq. ammonium chloride, dried and evaporated to a pale viscous oil. Ethyl acetate (300 ml) was added, the precipitate was filtered off and discarded, and the solution was
- 10 chromatographed (silica, 2.5%-10% methanol + 0.5% aq. ammonia in dichloromethane). Product


fractions were evaporated to an orange solid which solidified on standing (140 g). $^1\text{H-NMR}$ (d_6 -DMSO) δ 1.36 (9H,s), 3.37 (2H,s), 3.76 (3H,s), 4.42 (2H,s), 4.65 (2H,s), 7.13 (2H,m), 7.23 (1H,m), 7.45 (4H,m); MS (APCI+) found $(\text{M}+1) = 445$; $\text{C}_{22}\text{H}_{25}\text{FN}_4\text{O}_3\text{S}$ requires 444.

Intermediate B71 – 1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-((1-methylpyrazol-4-yl)methyl)-pyrimidin-4-one




1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-((1-methylpyrazol-4-yl)methyl)pyrimidin-4-one (96.8 g) was dissolved in dichloromethane (195 ml), cooled in ice/water, and trifluoroacetic acid (130 ml) added slowly with rapid stirring. After a further 36 hours stirring, the solvent was evaporated and the glassy residue triturated with ether; yield 78.6g. $^1\text{H-NMR}$ (d_6 -DMSO) δ 3.36 (2H, s), 3.76 (3H, s), 4.41 (2H, s), 4.67 (2H, s), 7.14 (2H, m), 7.23 (1H, s), 7.43-7.49 (4H, m); MS (APCI+) found $(\text{M}+1) = 389$; $\text{C}_{18}\text{H}_{17}\text{FN}_4\text{O}_3\text{S}$ requires 388.

Intermediate B72 – 1-Ethoxycarbonylmethyl-2-(4-fluorobenzyl)thio-5-(pyrimid-5-ylmethyl)-pyrimidin-4-one




A mixture of intermediate B52 (10g), ethyl bromoacetate (3.38ml), diisopropylethylamine (5.84ml) and dichloromethane (50ml) was stirred overnight, then the solution was washed sequentially with aqueous ammonium chloride and aqueous sodium bicarbonate. Chromatography (silica, 5-10% methanol in ethyl acetate) and trituration with ether gave the desired product (7.02g). $^1\text{H-NMR}$ (CDCl_3) δ 1.26 (3H, t), 3.71 (2H, s), 4.26 (2H, q), 4.46 (2H, s), 4.48 (2H, s), 6.91 (1H, s), 6.98 (2H, m), 7.35 (2H, m), 8.70 (2H, s), 9.09 (1H, s); MS(APCI $^+$) $\text{M}+1=415$, $\text{C}_{20}\text{H}_{19}\text{FN}_4\text{O}_3\text{S}$ requires 414. MPt 145.1°C.

Intermediate B73 – 1-Carboxymethyl-2-(4-fluorobenzyl)thio-5-(pyrimid-5-ylmethyl)pyrimidin-4-one




0.5M aqueous sodium hydroxide (33.8ml) was added slowly to a solution of Intermediate B72 (7.01g) in dioxan (150ml). The mixture was stirred for 2.5h at room temperature, then the dioxan was evaporated, water added, and the mixture acidified with aqueous sodium bisulfate. The precipitate was filtered off, washed with water and dried to obtain the desired product (6.31g). $^1\text{H-NMR}$ (d_6 -DMSO) δ 3.59 (2H, s), 4.41 (2H, s), 4.67 (2H, s), 7.11 (2H, m), 7.45 (2H, m), 7.72 (1H, s), 8.70 (2H, s), 9.03 (1H, s), 13.55 (1H, bs); MS (APCI-) $\text{M}-1=385$, $\text{C}_{18}\text{H}_{15}\text{FN}_4\text{O}_3\text{S}$ requires 386. MPt 206-207°C.

Intermediate B74 — 1-Ethoxycarbonylmethyl-2-(4-fluorobenzyl)thio-5-(2-oxopyrimidin-5-ylmethyl)pyrimidin-4-one



Prepared from Intermediate B86 by the method of Example 21, except using only 2 equivalents of B-bromocatecholborane. ¹H-NMR (d6-DMSO) δ 1.28 (3H, t), 4.26 (4H, q), 4.53 (2H, s), 4.90 (2H, s), 7.26 (2H, m), 7.57 (2H, m), 7.69 (1H, s), 8.25 (2H, br. s); MS (APCI+) found (M+1) = 431; C₂₀H₁₉FN₄O₄S requires 430.

Intermediate B75 — 1-Ethoxycarbonylmethyl-2-(4-fluorobenzyl)thio-5-(1-ethyl-2-oxopyrimidin-5-ylmethyl)pyrimidin-4-one



To a solution of Intermediate B74 (3.1g) in dry dimethylformamide (40ml) under argon was added ethyl iodide (1.4g) and anhydrous potassium carbonate (2.5g). The mixture was stirred at room temperature for 20h and the solvent removed *in vacuo*. The residue was partitioned between ethyl acetate and water. The organic layer was washed with brine and added in equal portions to three 10g silica cartridges. Elution of each column with EtOAc to 18% MeOH:EtOAc and combining appropriate fractions gave the title compound (1.5g). ¹H-NMR (CDCl₃) δ 1.27t1.39t), 3.46 (2H,s), 3.94 (2H,q), 4.25 (2H,q), 4.48 (2H,s), 4.57 (2H,s), 6.9-7.15 (2H,m), 7.21 (1H,s), 7.3-7.4 (2H,m), 7.9 (1H,m), 8.4 (1H,m), MS (APCI+) found (M+1) = 459; C₂₂H₂₃FN₄O₄S requires 458.

The following intermediates were prepared by the method of intermediate B70:

No	Precursors	Structure	Name
B7 6	Int. B57		1-(tert-Butoxycarbonylmethyl)-2-(3,4-difluorobenzyl)thio-5-(2-methoxypyrimidin-5-ylmethyl)pyrimidin-4-one
B7 7	Int. B58		1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-((1-(2-methoxyethyl)pyrazol-4-yl)methyl)pyrimidin-4-one
B7 8	Int. B60		1-(tert-Butoxycarbonylmethyl)-2-(2,6-dimethylpyridin-4-ylmethylthio)-5-((1-methylpyrazol-4-yl)methyl)pyrimidin-4-one

B7 9	Int. B64		1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-benzylpyrimidin-4-one
B8 0	Int. B65		1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-(4-chlorobenzyl)pyrimidin-4-one
B8 1	Int. B59		1-(tert-Butoxycarbonylmethyl)-2-benzylthio-5-((1-methylpyrazol-4-yl)methyl)pyrimidin-4-one
B8 2	Int. B66		1-(tert-Butoxycarbonylmethyl)-2-(3,4-difluorobenzylthio)-5-((1-methylpyrazol-4-yl)methyl)pyrimidin-4-one

The following intermediates were prepared by the method of intermediate B72:

No	Precursors	Structure	Name
B8 4	Int. B61		1-(Ethoxycarbonylmethyl)-2-(2,6-dichloropyrid-4-ylmethylthio)-5-(2-methoxypyrimid-5-ylmethyl)pyrimidin-4-one
B8 5	Int. B54		1-((Ethoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-(2-trifluoromethylpyrimid-5-ylmethyl)pyrimidin-4-one
B8 6	Int. B53		1-(Ethoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-(2-methoxypyrimid-5-ylmethyl)pyrimidin-4-one
B8 7	Int. B55		1-(Ethoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-(2-(4-morpholino)pyrimid-5-ylmethyl)pyrimidin-4-one
B8 8	Int. B56		1-(Ethoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-(2-dimethylaminopyrimid-5-ylmethyl)pyrimidin-4-one

The following intermediates were prepared by the method of intermediate B71:


No	Precurs rs	Structure	Name
B89	Int. B81		1-(Carboxymethyl)-2-benzylthio-5-(1-methylpyrazol-4-ylmethyl)pyrimidin-4-one
B90	Int. B76		1-(Carboxymethyl)-2-(3,4-difluorobenzyl)-thio-5-(2-methoxypyrimid-5-ylmethyl)-pyrimidin-4-one
B91	Int. B82		1-(Carboxymethyl)-2-(3,4-difluorobenzyl)-thio-5-(1-methylpyrazol-4-ylmethyl)-pyrimidin-4-one
B92	Int. B78		1-(Carboxymethyl)-2-(2,6-dimethylpyrid-4-ylmethylthio)-5-((1-methylpyrazol-4-yl)-methyl)pyrimidin-4-one
B93	Int. B79		1-Carboxymethyl-2-(4-fluorobenzylthio)-5-benzylpyrimidin-4-one
B94	Int. B80		1-Carboxymethyl-2-(4-fluorobenzylthio)-5-(4-chlorobenzyl)pyrimidin-4-one
B104	Int. B77		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-((1-(2-methoxyethyl)pyrazol-4-yl)methyl)pyrimidin-4-one

The following intermediates were prepared by the method of intermediate B73:

No.	Precursor s	Structure	Name
B96	Int. B51		1-(3-Carboxyprop-1-yl)-2-(4-fluorobenzyl)-thio-5-(2-methoxypyrimid-5-ylmethyl)-pyrimidin-4-one
B97	Int. B67		1-(3-Carboxyprop-1-yl)-2-(4-fluorobenzyl)-thio-5-(1-methylpyrazol-4-ylmethyl)-pyrimidin-4-one

B98	Int. B85		1-Carboxymethyl-2-(4-fluorobenzylthio)-5-((2-trifluoromethylpyrimid-5-yl)methyl)pyrimidin-4-one
B99	Int. B68		1-(3-Carboxyprop-1-yl)-2-(4-fluorobenzylthio)-5-(2-(4-morpholino)pyrimid-5-ylmethyl)pyrimidin-4-one
B100	Int. B75		1-Carboxymethyl-2-(4-fluorobenzylthio)-5-(1-ethyl-2-oxopyrimid-5-ylmethyl)pyrimidin-4-one
B101	Int. B87		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-(2-(4-morpholino)pyrimid-5-ylmethyl)pyrimidin-4-one
B102	Int. B88		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-(2-dimethylaminopyrimid-5-ylmethyl)pyrimidin-4-one
B103	Int. B84		1-(Carboxymethyl)-2-(2,6-dichloropyrid-4-ylmethylthio)-5-(2-methoxypyrimid-5-ylmethyl)pyrimidin-4-one
B104	Int. B86		1-Carboxymethyl-2-(4-Fluorobenzylthio)-5-(2-methoxypyrimid-5-ylmethyl)pyrimidin-4-one


Example 1 — 1-(N-Methyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methylpyrazol-4-ylmethyl)pyrimidin-4-one



A mixture of Intermediate A2 (0.27g, 1 equiv), Intermediate B71 (0.45g, 1 equiv),

- 5 hydroxybenzotriazole (0.018g, 0.1 equiv), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.25g, 1.1 equiv) and dichloromethane (15ml) was stirred at room temperature overnight, then diluted with dichloromethane and washed with aqueous sodium bicarbonate. The organic layer was applied directly to a 10g silica cartridge, which was eluted with 0-10% methanol in ethyl acetate. Product fractions were evaporated to an oil, which was triturated with ether to obtain a white solid (0.39g). ¹H-NMR (d₆-DMSO) δ 2.95 and 3.08 (3H, 2Xs), 3.61(2H, m), 3.86 (3H, m), 4.46-4.61 (6H, m), 6.74 and 6.80 (1H, 2Xs), 6.91-6.99 (2H, m), 7.21-7.49 (12H, m); MS (APCI+) found (M+1) = 602; C₃₂H₂₉ClFN₅O₂S requires 601.
- 10


Example 2 — 1-(N-Methyl-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methylpyrazol-4-ylmethyl)pyrimidin-4-one



15


Prepared from Intermediates A45 and B71 by the method of Example 1. ¹H-NMR (DMSO) δ 2.96 (3H,s), 3.39 (2H,m), 3.77 (3H,s), 4.43 (2H, s), 4.57 (2H,s), 4.97 (2H,s), 7.12 (3H,m), 7.25 (1H,s), 7.34-7.48 (6H, m), 7.62-6.70 (2H,m), 8.84 (4H,m).; MS (APCI+) found (M+1) = 636; C₃₃H₂₉F₄N₅O₂S requires 635.

- 20 **Example 3 — 1-(N-(2-Dimethylaminoethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methylpyrazol-4-ylmethyl)pyrimidin-4-one hydrochloride**




- Prepared from Intermediates A61 and B71 by the method of Example 1. ¹H-NMR (CDCl₃) δ 2.95 (3H, s), 2.96(3H, s), 3.19 (2H, m), 3.58 (2H, s), 3.87 (3H, s), 3.91 (2H, m), 4.42 (2H, s), 4.68 (2H, s), 4.99 (2H, s), 6.88-6.93 (2H, m), 7.25-7.29 (6H, m), 7.36-7.45 (6H, m) 7.62 (1H, s); MS (APCI+) found (M+1) = 659; C₃₅H₃₆ClFN₆O₂S requires 658.
- 25

Example 4 — 1-(N-Methyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-(4-methylphenyl)pyrimidin-5-ylmethyl)pyrimidin-4-one




A solution of Intermediate A7 (0.23g, 1 equiv), Intermediate B55 (0.27g, 1 equiv) and diisopropylethylamine (0.10g, 1.2 equiv) in dry dichloromethane (6ml) was stirred at room temperature under argon for 20h, then diluted with more dichloromethane and washed successively with water and aqueous ammonium chloride. Drying and evaporation of the organic phase, followed by chromatography (silica, 0-5% methanol in ethyl acetate) gave the desired product as a pale solid (0.19g). ¹H-NMR (DMSO-d₆) δ: 2.95 (d, 3H), 3.42 (d, 2H), 3.64 (s, 8H), 4.40 (d, 2H), 4.60 (d, 2H), 4.95 (d, 2H), 7.1 (m, 2H), 7.30 (d, 2H), 7.35-7.6 (m, 6H), 7.65 (m, 3H), 8.30 (s, 2H). MS (APCI+) Found (M+1) = 685/687; C₃₆H₃₄ClFN₆O₃S requires 685.

Example 5 — 1-(N-(2-(dimethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride



Prepared from Intermediates A72 and B71 by the method of Example 1. ¹H-NMR (CDCl₃) δ 2.98 (3H,s), 2.99 (3H,s), 3.20 (2H,m), 3.60 (2H,s), 3.93 (5H,m), 4.42 (3H,s), 4.69(2H,s), 5.00 (2H,s), 6.89 (2H,m), 7.2-7.3 (4H,m), 7.47 (4H, m), 7.55 (2H,d), 7.70 (3H,m), 11.7 (1H, br s); MS (APCI+) found (M+1) = 693; C₃₆H₃₆F₄N₆O₂S requires 692.

Example 6 — 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one



A mixture of Intermediate A3 (82.7g, 1 equiv), B71 (101.3g, 1 equiv), hydroxybenzotriazole (39.9g, 1 equiv), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (100.0g, 2 equiv) and dichloromethane (1200ml) was stirred under argon at room temperature overnight, then aqueous sodium bicarbonate was added slowly with stirring. The organic layer was separated, the aqueous layer extracted twice more with dichloromethane, and the combined organic extracts dried over potassium carbonate and evaporated to a brown oil. Trituration with ether gave a solid, which was filtered off and purified by

chromatography on silica, eluting with methanol/dichloromethane then with methanolic ammonia/dichloromethane. Product fractions were evaporated to a yellow foam (98g). ¹H-NMR (CDCl₃, rotamer mixture) δ 0.9-1.0 (6H,m), δ 2.4-2.6 (6H,m), 3.23/3.52 (4H,2x t), 3.58/3.61 (4H, 2x s), 3.85 (3H,s), 4.46/4.53/4.64/4.82 (6H,4x s), 6.75/6.79 (1H,2x s), 6.9 (2H,m), 7.2-7.5 (12H,m); MS (APCI+) found (M+1) = 687/689; C₃₇H₄₀ClFN₆O₂S requires 686/688.

Example 7 — 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride

The free base from Example 6 (8.7g, 1 equiv) was dissolved in dichloromethane (50ml) and 1M hydrogen chloride in ether (1 equiv) was added dropwise under argon. The mixture was evaporated to ca. half volume and sonicated to obtain a clear solution, which was transferred to a syringe and added dropwise to ether (200ml) with vigorous stirring. The white solid was filtered off, washed with ether and dried in vacuo; yield 8.55g. ¹H-NMR (DMSO, ca. 2:1 rotamer mixture) δ 1.14-1.24 (6H,m), 3.1 (6H, m), 3.6 (2H,m), 3.76 (3H,s), 4.38/4.45 (2H,2x s), 4.61/4.70 (2H,2x s), 4.98/5.11 (2H,2x s), 6.75/6.79 (1H,2x s), 7.1-7.7 (15H,m), 10.15/10.75 (1H,2x br s); MS (APCI+) found (M+1) = 687/689; C₃₇H₄₀ClFN₆O₂S requires 686/688.

Example 8 — 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one bitartrate

A solution of L-tartaric acid (0.75g, 1 equiv) in 2-propanol (12ml) was added to a solution of the free base from Example 6 (3.43g, 1 equiv) in 2-propanol (25ml) with stirring. The mixture was evaporated to ca. one third volume, diluted with ether, then the solid was filtered off, washed with ether and dried in vacuo; yield 3.87g. ¹H-NMR (DMSO, ca. 60:40 rotamer mixture) δ 0.9-1.1 (6H,m), 2.5-2.8 (4H,m), 3.2-3.4 (6H,m), 3.76 (3H,s), 4.20 (2H,s), 4.39/4.43 (2H,2x s), 4.60/4.67 (2H,2x s), 4.92/5.08 (2H,2x s), 7.1-7.7 (15H,m); MS (APCI+) found (M+1) = 687/689; C₃₇H₄₀ClFN₆O₂S requires 686/688.

Example 9 — 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one maleate

A solution of maleic acid (0.58g, 1 equiv) in methanol (10ml) was added to a solution of the free base from Example 6 (3.43g, 1 equiv) in methanol (10ml) with stirring. The mixture was evaporated to ca. half volume, diluted with ether, then the supernatant decanted off. The oil was triturated with ether to obtain a solid, which was filtered off, washed with ether and dried in vacuo; yield 3.69g. ¹H-NMR (DMSO, ca. 3:1 rotamer mixture) essentially similar to Example 7, plus δ 2.33 (3H,s); MS (APCI+) found (M+1) = 687/689; C₃₇H₄₀ClFN₆O₂S requires 686/688.


Example 10 — 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one mesylate

Prepared by the method of example 9, using methanesulfonic acid (0.32ml, 1 equiv) in place of maleic acid; yield 3.59g. ¹H-NMR (DMSO, ca. 3:1 rotamer mixture) essentially similar to Example 7, plus δ 6.03 (2H,s); MS (APCI+) found (M+1) = 687/689; C₃₇H₄₀ClFN₆O₂S requires 686/688.

Example 11 — 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one

Prepared by the method of example 9, using taurocholic acid (2.57g, 1 equiv) in place of maleic acid; yield 5.56g. ¹H-NMR (DMSO, ca. 3:1 rotamer mixture) essentially similar to Example 7, plus (*inter alia*) δ 0.58 (3H,s), 0.81 (3H,s), 0.91 (3H,d), 1.14 (3H,s); MS (APCI+) found (M+1) = 687/689; C₃₇H₄₀ClFN₆O₂S requires 686/688.

Example 12 — 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one



- 10 Prepared from Intermediates A73 and B71 by the method of Example 1. ¹H-NMR (CDCl₃, rotamer mixture) δ 0.9-1.0 (6H,m), δ 2.4-2.6 (6H,m), 3.24/3.4-3.6 (4H,2x m), 3.85 (3H,s), 4.46/4.53/4.66/4.83 (6H,4x s), 6.75/6.8 (1H,2x s), 6.9-7.0 (2H,m), 7.3-7.7 (12H,m); MS (APCI+) found (M+1) = 721; C₃₈H₄₀F₄N₆O₂S requires 720.

- 15 **Example 13 — 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride**

- 20 The free base from Example 12 (0.75g, 1 equiv) was dissolved in dichloromethane (5 ml) and 1M hydrogen chloride in ether (1 equiv) was added dropwise under argon, then excess ether was added with vigorous stirring. The white solid was filtered off, washed with ether and dried in vacuo; yield 0.73g. ¹H-NMR (DMSO, ca. 3:1 rotamer mixture) δ 1.1-1.2 (6H,m), 3.1 (6H, m), 3.37 (2H+H₂O,m), 3.66 (2H,m), 3.76 (3H,s), 4.37/4.45 (2H,2x s), 4.63/4.72 (2H,2x s), 4.97/5.12 (2H,2x s), 7.1 (2H,m), 7.24/7.26 (1H,2x s), 7.4-7.5 (6H,m), 6.62/6.71 (2H, 2x d), 7.84 (4H,m), 10.1/10.65 (1H,2x br s); MS (APCI+) found (M+1) = 721; C₃₈H₄₀F₄N₆O₂S requires 720.

- 25 **Example 14 — 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one bitartrate**

- 30 Prepared from the free base of Example 12 by the method of Example 8. ¹H-NMR (DMSO, ca. 60:40 rotamer mixture) δ 0.9-1.0 (6H,m), 2.5-2.8 (4H,m), 3.2-3.5 (6H,m), 3.76 (3H,s), 4.19 (2H,s), 4.38/4.43 (2H,2x s), 4.62/4.69 (2H,2x s), 4.92/5.10 (2H,2x s), 7.0-7.5 (12H,m), 7.6/7.7 (1H, 2x d), 7.84 (2H,m); MS (APCI+) found (M+1) = 721; C₃₈H₄₀F₄N₆O₂S requires 720.


Example 15 — 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one citrate

Free base from Example 12 (0.69g, 1equiv) was added to acetone (10ml) followed by citric acid (0.20g, 1equiv). The mixture was warmed to dissolve the solids then allowed to cool, whereupon the solids were filtered off and dried in vacuo; yield 0.80g, 90%, m.p. 130-133°C. ¹H-NMR (d6-DMSO, ca. 1:1 rotamer mixture) essentially similar to Example 14, plus δ 2.61 (2H,d); MS (APCI+) found (M+1) = 721; C₃₈H₄₀F₄N₆O₂S requires 720.

Example 16 — 1-(N-(2-(diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one tosylate

Free base from Example 12 (1.0 g, 1 equiv) and tosic acid (0.26 g, 1 equiv) were dissolved in acetone (20 ml) at room temperature, then isopropyl acetate (60 ml) was added with stirring. After stirring for a further 2 hours, the mixture was allowed to stand overnight then the solid was filtered off, washed with 3:1 isopropyl acetate/acetone, and dried in vacuo; yield 0.87 g, m.p. 147°C. ¹H-NMR (d6-DMSO, ca. 3:1 rotamer mixture) essentially similar to Example 13, plus (inter alia) δ 2.28 (3H,s); MS (APCI+) found (M+1) = 721; C₃₈H₄₀F₄N₆O₂S requires 720

Example 17 — 1-(N-(2-(diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-ylmethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one



A mixture of Intermediate A106 (0.523g, 1 equiv), Intermediate B71 (0.473g, 1 equiv), HATU (0.55g, 1.2 equiv), diisopropylethylamine (0.622ml, 2.4 equiv) and dichloromethane (20ml) was stirred under argon at room temperature overnight, then washed with aqueous ammonium chloride and aqueous sodium bicarbonate. The organic layer was dried and evaporated, and the free-base product isolated by chromatography (silica, 3-8% methanol in dichloromethane) as a pale foam (0.39g). ¹H-NMR (CDCl₃, major rotamer) δ 0.94 (6H,t), 2.47 (4H,q), 2.58 (2H,m), 3.26 (2H,m), 3.61 (2H,s), 3.85 (3H,s), 4.52 (2H,s), 4.66 (2H,s), 4.82 (2H,s), 6.78 (1H,s), 6.98 (2H,m), 7.3-7.4 (4H,m), 7.6-7.8 (2H,m), 8.08 (2H,m), 8.57 (1H,m); MS (APCI+) found (M+1) = 722; C₃₇H₃₉F₄N₇O₂S requires 721.

Example 18 — 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-ylmethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride


The free base from Example 17 (0.373g, 1 equiv) was dissolved in dichloromethane (10ml), and a solution of hydrogen chloride in ether (0.517ml, 1.0M solution, 1 equiv) was added dropwise with stirring. The solvent was removed in vacuo, and the residue triturated with ether to obtain a white solid (0.362g). ¹H-NMR (DMSO, ca. 60:40 rotamer mixture) δ 1.1-1.3 (6H,m), 3.77 (3H,s), 4.38/4.43 (2H,s), 4.65/4.76 (2H,s), 5.00/5.08 (2H,s), 7.10 (2H,m), 7.25 (1H,m), 7.3-7.5 (4H,m), 7.78 (1H,m), 7.87

(2H,d), 7.97/8.06 (1H,d), 8.28 (2H,d), 8.61/8.70 (1H,m); MS (APCI+) found (M+1) = 722;
 $C_{37}H_{39}F_4N_7O_2S$ requires 721.

Example 19 — 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-ylmethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one bitartrate


Prepared from the free base of Example 17 by the method of Example 8. 1H -NMR (DMSO, ca. 60:40 rotamer mixture) δ 0.88-1.05 (6H,m), 3.76 (3H,s), 4.39/4.42 (2H,s), 4.65/4.74 (2H,s), 4.97/5.10 (2H,s), 7.11 (2H,m), 7.25 (1H,m), 7.3-7.5 (4H,m), 7.78 (1H,m), 7.87 (2H,m), 7.98/8.07 (1H,d), 8.28 (2H,d), 8.61/8.70 (1H,m); MS (APCI+) found (M+1) = 722; $C_{37}H_{39}F_4N_7O_2S$ requires 721.

Example 20 — 1-(N-(2-(1-Piperidino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one bitartrate




The free base was prepared from Intermediates A108 and B71 by the method of Example 1, then converted to bitartrate salt by the method of Example 8. 1H -NMR (DMSO, ca. 60:40 rotamer mixture) δ 1.3-1.6 (6H,m), δ 2.3-2.7 (6H,m), 3.3-3.6 (4H,m), 3.77 (3H,s), 4.22 (2H,s), 4.37/4.44 (2H,2x s), 4.62/4.69 (2H,2x s), 4.88/5.08 (2H,2x s), 7.1-7.5 (12H,m), 7.6/7.7 (1H, 2x d), 7.83 (2H,m); MS (APCI+) found (M+1) = 733; $C_{39}H_{40}F_4N_6O_2S$ requires 732.

Example 21 — 1-(N-(Carboxymethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one sodium salt




Example 102 (130 mg) was added to a solution of sodium bicarbonate (16 mg) in water (4 ml) and stirred for 1 hour at room temperature. A small proportion of methanol was added to obtain a clear solution on warming and sonication. Filtration and lyophilisation gave the desired sodium salt as a white solid. 1H -NMR (DMSO) δ 3.49 (2H,s), 3.76 (3H,s), 4.42 (2H,s), 4.85 (2H,s), 7.14 (2H,m), 7.26 (1H,s), 7.36 (3H,m), 7.49 (3H,m), 7.60 (2H,m), 7.84 (4H,m); MS (APCI-) found (M+1) = 678; $C_{34}H_{29}F_4N_5O_4S$ requires 679.

Example 22 — 1-(N-(2-aminoethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride



The Boc-protected product was prepared from Intermediates A84 and B71 by the method of Example 1, then deprotected by suspending in dioxan and treating with excess hydrogen chloride (4M solution in dioxan). After stirring for 1 hour, excess ether was added and stirring continued until a fine white solid was obtained. This was filtered off, washed with ether and dried to obtain the crude product., which was purified by chromatography (silica, 10-20% methanol in dichloromethane). ¹H-NMR (DMSO, ca. 60:40 rotamer mixture) δ 2.97/3.04 (2H,m) δ 3.37 (2H,m), 3.55 (2H,m), 3.77 (3H,s), 4.36/4.45 (2H,2x s), 4.60/4.67 (2H,2x s), 4.93/5.09 (2H,2x s), 7.0-7.7 (15H,m), 8.05/8.20 (2H,2x br s); MS (APCI+) found (M+1) = 733; C₃₉H₄₀F₄N₆O₂S requires 732.

Example 23 — 1-(N-Methyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-oxopyrimidin-5-ylmethyl)pyrimidin-4-one












A mixture of Example 121 (0.73g, 1.2mmol), B-bromocatecholborane (1.01g, 5.1mmol) and dry dichloromethane (20ml) was stirred under argon at room temperature overnight, giving a clear orange solution. Water was added and stirring continued for 30min, then the organic layer was separated and applied directly to a 10g silica cartridge, which was eluted with 0-14% methanol in dichloromethane. Product fractions were evaporated to obtain a pale yellow solid (0.36g). ¹H-NMR (DMSO) δ 2.80-3.06 (3H, 2xs), 3.37 (2H, s), 4.33-4.49 (2H, 2xs), 4.50-4.72 (2H, 2xs), 4.87-5.06 (2H, m), 7.04-7.75 (13H, m), 8.19 (2H, s); MS (APCI-) found (M-1) = 614; C₃₂H₂₇N₅O₃SFCl requires 615.

The following Examples were made either by the method of Example 1 (EDC coupling) or Example 15 (HATU coupling); where indicated, the salts were subsequently prepared by the methods of Examples 7-8 as appropriate:

Ex No	Precursors	Structure	Name

30	Int. A2 Int. B97		1-(3-(N-methyl-N-(4-(4-chlorophenyl)-benzyl)aminocarbonyl)prop-1-yl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
31	Int. A120 Int. B97		1-(3-(4-(4-fluorophenyl)benzylamino-carbonyl)prop-1-yl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolyl-methyl)pyrimidin-4-one
32	Int. A44 Int. B97		1-(3-(N-methyl-N-(4-(4-fluorophenyl)-benzyl)aminocarbonyl)prop-1-yl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
33	Int. A124 Int. B97		1-(3-(4-phenylbenzylaminocarbonyl)-prop-1-yl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
34	Int. A6 Int. B99		1-(3-(4-(4-chlorophenyl)benzylamino-carbonyl)prop-1-yl)-2-(4-fluorobenzyl)-thio-5-(2-(4-morpholino)-5-pyrimidyl-methyl)pyrimidin-4-one
35	Int. A2 Int. B99		1-(3-(N-methyl-N-(4-(4-chlorophenyl)-benzyl)aminocarbonyl)prop-1-yl)-2-(4-fluorobenzyl)thio-5-(2-(4-morpholino)-5-pyrimidylmethyl)pyrimidin-4-one
36	Int. A6 Int. B96		1-(3-(4-(4-chlorophenyl)benzylamino-carbonyl)prop-1-yl)-2-(4-fluorobenzyl)-thio-5-(2-methoxy-5-pyrimidylmethyl)-pyrimidin-4-one
37	Int. A2 Int. B96		1-(3-(N-methyl-N-(4-(4-chlorophenyl)-benzyl)aminocarbonyl)prop-1-yl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one

38	Int. A120 Int. B96		1-(3-(4-(4-fluorophenyl)benzylamino-carbonyl)prop-1-yl)-2-(4-fluorobenzyl)-thio-5-(2-methoxy-5-pyrimidylmethyl)-pyrimidin-4-one
39	Int. A44 Int. B96		1-(3-(N-methyl-N-(4-(4-fluorophenyl)-benzyl)aminocarbonyl)prop-1-yl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
40	Int. A124 Int. B96		1-(3-(4-phenylbenzylaminocarbonyl)-prop-1-yl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
41	Int. A64 Int. B96		1-(3-(N-methyl-N-(2-(4-fluorophenyl)-pyrid-5-ylmethyl)aminocarbonyl)prop-1-yl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
42	Int. A3 Int. B104		1-(N-(2-(diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-(1-(2-methoxyethyl)-4-pyrazolylmethyl)-pyrimidin-4-one bitartrate
43	Int. A73 Int. B104		1-(N-(2-(diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-(2-methoxyethyl)-4-pyrazolyl-methyl)pyrimidin-4-one bitartrate
44	Int. A6 Int. B100		1-(4-(4-chlorophenyl)benzylamino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-ethyl-2-oxo-5-pyrimidylmethyl)-pyrimidin-4-one
45	Int. A2 Int. B100		1-(N-methyl-N-(4-(4-chlorophenyl)-benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-ethyl-2-oxo-5-pyrimidylmethyl)pyrimidin-4-one
46	Int. A44 Int. B100		1-(N-methyl-N-(4-(4-fluorophenyl)-benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-ethyl-2-oxo-5-pyrimidylmethyl)pyrimidin-4-one









47	Int. A2 Int. B92		1-(N-methyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(2,6-dimethylpyrid-4-yl)methylthio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
48	Int. A3 Int. B90		1-(N-(2-(diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(3,4-difluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
49	Int. A45 Int. B90		1-(N-methyl-N-(4-(4-(trifluoromethyl)phenyl)benzyl)aminocarbonylmethyl)-2-(3,4-difluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
50	Int. A46 Int. B71		1-(N-methyl-N-(4-(4-chlorophenyl)-2-fluorobenzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
51	Int. A56 Int. B71		1-(N-methyl-N-(4-(2,4-bis(trifluoromethyl)phenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
52	Int. A60 Int. B71		1-(N-methyl-N-(4-(2,4-difluorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
53	Int. A59 Int. B71		1-(N-methyl-N-(4-(3-chloro-4-fluorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
54	Int. A53 Int. B71		1-(N-methyl-N-(4-(3-fluoro-4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
55	Int. A107 Int. B71		1-(N-(2-(1-piperidino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride

56	Int. A75 Int. B71		1-(N-(2-(1-pyrrolidino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
57	Int. A76 Int. B71		1-(N-(2-(4-morpholino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
58	Int. A95 Int. B71		1-(N-(2-(bis(2-hydroxyethyl)amino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
59	Int. A74 Int. B71		1-(N-(2-(di-isopropylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
60	Int. A94 Int. B71		1-(N-(2-(N-(2-hydroxyethyl)-N-ethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one bitartrate
61	Int. A70 Int. B71		1-(N-(2-hydroxyethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
62	Int. A51 Int. B71		1-(N-(2-methoxyethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
63	Int. A80 Int. B71		1-(N-(3-(1-pyrrolidino)propyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride

64	Int. A83 Int. B71		1-(N-(3-(4-methyl-1-piperazino)propyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)-pyrimidin-4-one hydrochloride
65	Int. A82 Int. B71		1-(N-(3-(4-morpholino)propyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
66	Int. A79 Int. B71		1-(N-(3-(diethylamino)propyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
67	Int. A62 Int. B71		1-(N-(3-(dimethylamino)propyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
68	Int. A50 Int. B71		1-(N-(dimethylaminocarbonylmethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)-pyrimidin-4-one
69	Int. A48 Int. B71		1-(N-ethyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
70	Int. A6 Int. B71		1-(4-(4-chlorophenyl)benzylaminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)-pyrimidin-4-one
71	Int. A93 Int. B71		1-(N-(2-(diethylamino)ethyl)-N-(4-(4-difluoromethoxyphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride

72	Int. A120 Int. B71		1-(4-(4-fluorophenyl)benzylamino)carbonylmethyl-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
73	Int. A44 Int. B71		1-(N-methyl-N-(4-(4-fluorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
74	Int. A54 Int. B71		1-(N-methyl-N-(4-(4-methoxyphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
75	Int. A92 Int. B71		1-(N-(2-(diethylamino)ethyl)-N-(4-(4-trifluoromethoxyphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
76	Int. A71 Int. B71		1-(N-(2-hydroxyethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
77	Int. A78 Int. B71		1-(N-(3-(dimethylamino)propyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
78	Int. A97 Int. B71		1-(N-(4-methyl-1-piperazinocarbonylmethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
79	Int. A96 Int. B71		1-(N-(4-morpholinocarbonylmethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one

80	Int. A52 Int. B71		1-(N-(dimethylaminocarbonylmethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
81	Int. A121 Int. B71		1-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
82	Int. A98 Int. B71		1-(N-(2-(diethylamino)ethyl)-N-(5-(4-chlorophenyl)pyrid-2-ylmethyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)-pyrimidin-4-one hydrochloride
83	Int. A102 Int. B71		1-(N-(2-(dimethylamino)ethyl)-N-(5-(4-trifluoromethoxyphenyl)pyrid-2-yl-methyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
84	Int. A99 Int. B71		1-(N-(2-(diethylamino)ethyl)-N-(5-(4-trifluoromethylphenyl)pyrid-2-ylmethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
85	Int. A101 Int. B71		1-(N-(2-(diethylamino)ethyl)-N-(2-(4-chlorophenyl)pyrid-5-ylmethyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)-pyrimidin-4-one hydrochloride
86	Int. A103 Int. B71		1-(N-(2-hydroxyethyl)-N-(2-(4-chloro-phenyl)pyrid-5-ylmethyl)aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
87	Int. A122 Int. B71		1-(2-(4-fluorophenyl)pyrid-5-ylmethyl-aminocarbonylmethyl)-2-(4-fluoro-benzyl)thio-5-(1-methyl-4-pyrazolyl-methyl)pyrimidin-4-one

88	Int. A64 Int. B71		1-(N-methyl-N-(2-(4-fluorophenyl)pyrid-5-ylmethyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
89	Int. A100 Int. B71		1-(N-(2-(dimethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-ylmethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
90	Int. A104 Int. B71		1-(N-(2-(diethylamino)ethyl)-N-(2-(4-chlorophenyl)pyrimid-5-ylmethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one bitartrate
91	Int. A123 Int. B71		1-(5-(4-chlorophenyl)thien-2-ylmethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
92	Int. A66 Int. B71		1-(N-methyl-N-(5-(4-chlorophenyl)thien-2-ylmethyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
93	Int. A63 Int. B71		1-(N-methyl-N-(5-(4-fluorophenyl)thien-2-ylmethyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
94	Int. A3 Int. B89		1-(N-(2-(diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-benzylthio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
95	Int. A83 Int. B101		1-(N-(3-(4-methyl-1-piperazino)propyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-(4-morpholino)-5-pyrimidylmethyl)pyrimidin-4-one bitartrate

96	Int. A45 Int. B101		1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-(4-morpholino)-5-pyrimidylmethyl)-pyrimidin-4-one
97	Int. A83 Int. B102		1-(N-(3-(4-methyl-1-piperazino)propyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-dimethylamino-5-pyrimidylmethyl)-pyrimidin-4-one bitartrate
98	Int. A2 Int. B102		1-(N-methyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-dimethylamino-5-pyrimidylmethyl)pyrimidin-4-one
99	Int. A45 Int. B102		1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-dimethylamino-5-pyrimidylmethyl)pyrimidin-4-one
100	Int. A73 Int. B103		1-(N-(2-(diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(2,6-dichloropyrid-4-yl)methylthio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one bitartrate
101	Int. A45 Int. B103		1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(2,6-dichloropyrid-4-yl)methylthio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
102	Int. A160 Int. B71		1-(N-(Ethoxycarbonylmethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one
103	Int. A2 Int. B90		1-(N-methyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(3,4-difluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one

10 4	Int. A45 Int. B90		1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(3,4-difluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
10 5	Int. A46 Int. B104		1-(N-methyl-N-(4-(4-chlorophenyl)-2-fluorobenzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
10 6	Int. A58 Int. B104		1-(N-methyl-N-(4-(4-trifluoromethylphenyl)-2-fluorobenzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
10 7	Int. A10 Int. B104		1-(N-methyl-N-(4-(4-chlorophenyl)-3-fluorobenzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
10 8	Int. A57 Int. B104		1-(N-methyl-N-(4-(2-fluoro-4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
10 9	Int. A60 Int. B104		1-(N-methyl-N-(4-(2,4-difluorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
11 0	Int. A55 Int. B104		1-(N-methyl-N-(4-(2-fluoro-4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
11 1	Int. A59 Int. B104		1-(N-methyl-N-(4-(3-chloro-4-fluorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
11 2	Int. A53 Int. B104		1-(N-methyl-N-(4-(3-fluoro-4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one

11 3	Int. A3 Int. B104		1-(N-(2-(diethylamin)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one hydrochloride
11 4	Int. A61 Int. B104		1-(N-(2-(dimethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one hydrochloride
11 5	Int. A70 Int. B104		1-(N-(2-hydroxyethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
11 6	Int. A51 Int. B104		1-(N-(2-methoxyethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
11 7	Int. A50 Int. B104		1-(N-(dimethylaminocarbonylmethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
11 8	Int. A11 Int. B104		1-(N-(ethoxycarbonylmethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
11 9	Int. A48 Int. B104		1-(N-ethyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
12 0	Int. A6 Int. B104		1-(4-(4-chlorophenyl)benzylamino)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
12 1	Int. A2 Int. B104		1-(N-methyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one

12 2	Int. A120 Int. B104		1-(4-(4-fluorophenyl)benzylamino)- carbonylmethyl)-2-(4-fluorobenzyl)thio- 5-(2-methoxy-5-pyrimidylmethyl)- pyrimidin-4-one
12 3	Int. A44 Int. B104		1-(N-methyl-N-(4-(4-fluorophenyl)- benzyl)aminocarbonylmethyl)-2-(4- fluorobenzyl)thio-5-(2-methoxy-5- pyrimidylmethyl)pyrimidin-4-one
12 5	Int. A52 Int. B104		1-(N-(dimethylaminocarbonylmethyl)-N- (4-(4-trifluoromethylphenyl)benzyl)- aminocarbonylmethyl)-2-(4-fluoro- benzyl)thio-5-(2-methoxy-5-pyrimidyl- methyl)pyrimidin-4-one
12 6	Int. A45 Int. B104		1-(N-methyl-N-(4-(4-trifluoromethyl- phenyl)benzyl)aminocarbonylmethyl)-2- (4-fluorobenzyl)thio-5-(2-methoxy-5- pyrimidylmethyl)pyrimidin-4-one
12 7	Int. A49 Int. B104		1-(N-methyl-N-(2-(3,4-dichlorophenyl)- pyrid-5-ylmethyl)aminocarbonylmethyl)- 2-(4-fluorobenzyl)thio-5-(2-methoxy-5- pyrimidylmethyl)pyrimidin-4-one
12 8	Int. A65 Int. B104		1-(2-(4-chlorophenyl)pyrid-5-ylmethyl)- aminocarbonylmethyl)-2-(4-fluoro- benzyl)thio-5-(2-methoxy-5-pyrimidyl- methyl)pyrimidin-4-one
13 0	Int. A6 Int. B98		1-(4-(4-chlorophenyl)benzylamino)- carbonylmethyl)-2-(4-fluorobenzyl)thio- 5-(2-trifluoromethyl-5-pyrimidylmethyl)- pyrimidin-4-one
13 1	Int. A2 Int. B98		1-(N-methyl-N-(4-(4-chlorophenyl)- benzyl)aminocarbonylmethyl)-2-(4- fluorobenzyl)thio-5-(2-trifluoromethyl-5- pyrimidylmethyl)pyrimidin-4-one
13 2	Int. A73 Int. B94		1-(N-(2-(diethylamino)ethyl)-N-(4-(4- trifluoromethylphenyl)benzyl)amino- carbonylmethyl)-2-(4-fluorobenzyl)thio- 5-(4-chlorophenylmethyl)pyrimidin-4- one bitartrate

13 3	Int. A45 Int. B94		1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(4-chlorophenylmethyl)pyrimidin-4-one
13 4	Int. A124 Int. B73		1-(4-phenylbenzylaminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(5-pyrimidylmethyl)pyrimidin-4-one
13 5	Int. A67 Int. B73		1-(N-methyl-N-(4-phenylbenzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(5-pyrimidylmethyl)pyrimidin-4-one
13 6	Int. A73 Int. B93		1-(N-(2-(diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(phenylmethyl)pyrimidin-4-one bitartrate
13 7	Int. A45 Int. B93		1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(phenylmethyl)pyrimidin-4-one
13 8	Int. A73 Int. B101		1-(N-(2-(diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-(4-morpholino)pyrimid-5-ylmethyl)pyrimidin-4-one


The following Examples were made by the method of Example 4; the salts were subsequently prepared by the methods of Examples 7-8 as appropriate:

Ex. No.	Precursors	Structure	Name
140	Int. A155 Int. B63		1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(2-methylthiazol-4-yl)methylthio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one
141	Int. A155 Int. B62		1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(pyrid-3-yl)methylthio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one

The following Examples were made by the method of Example 22; the salts were subsequently prepared by the methods of Examples 7-8 as appropriate:


Ex. No.	Precursors	Structure	Name
150	Int. A90 Int. B71		1-(N-(2-(1-piperazino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
151	Int. A86 Int. B71		1-(N-(2-(ethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
152	Int. A85 Int. B71		1-(N-(2-(methylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
153	Int. A91 Int. B71		1-(N-(3-(2-oxo-1-piperazino)propyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
154	Int. A89 Int. B71		1-(N-(3-(methylamino)propyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
155	Int. A88 Int. B71		1-(N-(3-aminopropyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one hydrochloride
156	Int. A87 Int. B71		1-(N-(2-(ethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(1-methyl-4-pyrazolylmethyl)pyrimidin-4-one bitartrate

The following Example was made by the method of Example 21:

Ex. No.	Precurs or	Structure	Name
157	Ex. 118		1-(N-(carboxymethyl)-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methoxy-5-pyrimidylmethyl)pyrimidin-4-one, sodium salt

Biological Data**1. Screen for Lp-PLA₂ inhibition.**

- 5 Enzyme activity was determined by measuring the rate of turnover of the artificial substrate (A) at 37 °C in 50 mM HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulphonic acid) buffer containing 150 mM NaCl, pH 7.4.



10

(A)

Assays were performed in 96 well titre plates.

- Recombinant Lp PLA₂ was purified to homogeneity from baculovirus infected Sf9 cells, using a zinc chelating column, blue sepharose affinity chromatography and an anion exchange column. Following
- 15 purification and ultrafiltration, the enzyme was stored at 6 mg/ml at 4°C. Assay plates of compound or vehicle plus buffer were set up using automated robotics to a volume of 170 µl. The reaction was initiated by the addition of 20µl of 10x substrate (A) to give a final substrate concentration of 20 µM and 10 µl of diluted enzyme to a final 0.2 nM LpPLA₂.

- 20 The reaction was followed at 405 nm and 37 °C for 20 minutes using a plate reader with automatic mixing. The rate of reaction was measured as the rate of change of absorbance.

Results

- 25 The compounds described in the Examples were tested as described above and had IC₅₀ values in the range 0.001 to 0.00005 µM.